




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/415, C12N 15/82, A01H 5/00		A1	(11) International Publication Number: WO 00/01722
			(43) International Publication Date: 13 January 2000 (13.01.00)
(21) International Application Number: PCT/US99/15255 (22) International Filing Date: 7 July 1999 (07.07.99) (30) Priority Data: 60/091,875 7 July 1998 (07.07.98) US (71) Applicant (for all designated States except US): PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BRIGGS, Steven, P. [US/US]; 2454 Mango Way, Del Mar, CA 92014 (US). SIMMONS, Carl, R. [US/US]; 4228 Holland Drive, Des Moines, IA 50310 (US). (74) Agents: SPRUILL, W., Murray et al.; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: MANIPULATION OF MLO GENES TO ENHANCE DISEASE RESISTANCE IN PLANTS

Maize MLO Resistance Gene Homologues



(57) Abstract

Compositions and methods for enhancing disease resistance in plants are provided. The method involves modulating the activity of *Mlo* sequences in the plant. Particular *Mlo* sequences are provided that can be manipulated to enhance pathogen resistance in modified plants. Transformed plants, plant cells, tissues, and seed are also provided having enhanced disease resistance.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MANIPULATION OF *MLO* GENES TO ENHANCE DISEASE RESISTANCE IN PLANTS

5 FIELD OF THE INVENTION

The invention relates to the genetic manipulation of plants, particularly to transforming plants with genes that enhance disease resistance.

BACKGROUND OF THE INVENTION

10 Disease in plants is caused by biotic and abiotic causes. Biotic causes include fungi, viruses, bacteria, and nematodes. Of these, fungi are the most frequent causative agent of disease on plants. Abiotic causes of disease in plants include extremes of temperature, water, oxygen, soil pH, plus nutrient-element deficiencies and imbalances, excess heavy metals, and air pollution.

15 A host of cellular processes enables plants to defend themselves from disease caused by pathogenic agents. These processes apparently form an integrated set of resistance mechanisms that is activated by initial infection and then limits further spread of the invading pathogenic microorganism. This limitation of the pathogen intruder is frequently accomplished by localized
20 containment of the intruder following a coordinated complex defense response.

Subsequent to recognition of a potentially pathogenic microbe, plants can activate an array of biochemical responses. Generally, the plant responds by inducing several local responses in the cells immediately surrounding the infection site. The most common resistance response observed in both nonhost
25 and race-specific interactions is termed the "hypersensitive response" (HR). In the hypersensitive response, cells contacted by the pathogen, and often neighboring cells, rapidly collapse and dry in a necrotic fleck. Other responses include the deposition of callose, the physical thickening of cell walls by lignification, and the synthesis of various antibiotic small molecules and
30 proteins. Genetic factors in both the host and the pathogen determine the specificity of these local responses, which can be very effective in limiting the spread of infection.

The hypersensitive response in many plant-pathogen interactions is specified by and dependent on the presence of two complementary genes, one from the host and one from the pathogen. These complementary genes are the resistance (*R*) gene in the plant and a corresponding avirulence (*avr*) gene in the pathogen. The interaction of the genes is associated with the rapid, localized cell death of the hypersensitive response. *R* genes that respond to specific bacterial, fungal, or viral pathogens, have been isolated from a variety of plant species and several appear to encode cytoplasmic proteins.

The resistance gene in the plant and the avirulence gene in the pathogen often conform to a gene-for-gene relationship. That is, resistance to a pathogen is only observed when the pathogen carries a specific avirulence gene and the plant carries a corresponding or complementing resistance gene. Because *avr-R* gene-for-gene relationships are observed in many plant-pathogen systems and are accompanied by a characteristic set of defense responses, a common molecular mechanism underlying *avr-R* gene mediated resistance has been postulated. A simple model which has been proposed is that pathogen *avr* genes directly or indirectly generate a specific molecular signal (ligand) that is recognized by cognate receptors encoded by plant *R* genes.

Both plant resistance genes and corresponding pathogen avirulence genes have been cloned. The plant kingdom contains thousands of *R* genes with specific specificities for viral, bacterial, fungal, or nematode pathogens. Although there are differences in the defense responses induced during different plant-pathogen interactions, some common themes are apparent among *R* gene-mediated defenses. The function of a given *R* gene is dependent on the genotype of the pathogen. Plant pathogens produce a diversity of potential signals, and in a fashion analogous to the production of antigens by mammalian pathogens, some of these signals are detectable by some plants.

The avirulence gene causes the pathogen to produce a signal that triggers a strong defense response in a plant with the appropriate *R* gene. However, expressing an avirulence gene does not stop the pathogen from being virulent on hosts that lack the corresponding *R* gene. A single plant can have many *R* genes, and a pathogen can have many *avr* genes.

Monogenic resistance mediated by recessive (*mlo*) alleles of the *Mlo* locus is different. It differs from race-specific incompatibility to single pathogen strains in that it is believed to confer a broad spectrum resistance to almost all known isolates of the fungal pathogen, and the resistance is
5 apparently durable in the field despite extensive cultivation. Further, *mlo* resistance alleles have been obtained by mutagen treatment of susceptible wild-type *Mlo* varieties. These *mlo* plants exhibit a spontaneous leaf cell death phenotype under pathogen-free or even axenic conditions.

As noted, among the causative agents of infectious disease of crop
10 plants, the phytopathogenic fungi play the dominant role. Phytopathogenic fungi cause devastating epidemics, as well as causing significant annual crop yield losses. All of the approximately 300,000 species of flowering plants are attacked by pathogenic fungi. However, a single plant species can be host to only a few fungal species, and similarly, most fungi usually have a limited host
15 range.

Plant disease outbreaks have resulted in catastrophic crop failures that have triggered famines and caused major social change. Generally, the best strategy for plant disease control is to use resistant cultivars selected or developed by plant breeders for this purpose. However, the potential for serious
20 crop disease epidemics persists today, as evidenced by outbreaks of the Victoria blight of oats and southern corn leaf blight.

Accordingly, molecular methods are needed to supplement traditional breeding methods to protect plants from pathogen attack. Particularly, methods are needed for broad spectrum resistance to pathogens.

25

SUMMARY OF THE INVENTION

Compositions and methods for creating or enhancing resistance to plant pests are provided. The method provides control of pathogens by modulating the expression of *Mlo* genes. Novel *Mlo* sequences are provided from maize.

5 Such sequences can be utilized to modulate the expression of *Mlo* genes in plants, particularly maize, to enhance resistance to pathogens. Generally, such modulation will result in decreased or increased expression of native *Mlo* genes, preferably decreased expression. Such decreased expression can be effected by mutagenesis or expression of modified or antisense *Mlo* sequences described
10 herein.

It is recognized that a variety of promoters will be useful in the invention the choice of which will depend in part upon the desired level of expression of the modified sequences in the plant or alternatively, in the plant organ. It is recognized that the levels of expression can be controlled to induce
15 broad spectrum resistance resulting in levels of immunity in the plant or to induce cell death.

The methods of the invention find use in controlling plant pests, including fungal pathogens, viruses, nematodes, insects, and the like. Transformed plants and seeds, as well as methods for making such plants and
20 seeds are additionally provided.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a sequence alignment for the maize *Mlo* homologues with the barley *Mlo* sequence. The barley sequence is designated as Hv*Mlo*1.

25 Figure 2 groups the *Mlo* sequences based on relatedness as well as the plant tissue from which it was isolated.

Figure 3 schematically illustrates a plasmid vector comprising the Zm*Mlo*1 antisense construct operably linked to the ubiquitin promoter.

DETAILED DESCRIPTION OF THE INVENTION

Compositions of the invention include eight mutation-induced recessive alleles of maize set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13 and 15. The maize sequences exhibit homology to the *Mlo* barley sequence (SEQ ID NO: 17). See, Büschges *et al.* (1997) 88:695-705 and Tables 1 and 2. The isolated maize *Mlo* genes are involved in enhancing resistance to plant pests. In particular, the present invention provides for isolated nucleic acid molecules comprising nucleotide sequences encoding the amino acid sequences shown in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, or the nucleotide sequences encoding the DNA sequences deposited in a bacterial host with the American Type Culture Collection and assigned the Accession Nos. 98725, 98726, 98727, 98728, 98729 (SEQ ID NOS: 3, 5, 9, 11, 13 respectively). Further provided are polypeptides having an amino acid sequence encoded by a nucleic acid molecule described herein, for example those set forth in SEQ ID NOS: 1, 5, 3, 7, 9, 11, 13, and 15, those deposited with the American Type Culture Collection and assigned Accession Nos. 98725, 98726, 98727, 98728, 98729 (SEQ ID NOS: 3, 5, 9, 11, 13 respectively), and fragments and variants thereof.

Plasmids containing the nucleotide sequences of the invention were deposited with American Type Culture Collection, Manassas, Virginia, and assigned Accession Nos. 98725, 98726, 98727, 98728, 98729 (SEQ ID NOS: 3, 5, 9, 11, 13 respectively). These deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. These deposits were made merely as a convenience for those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112.

The invention encompasses isolated or substantially purified nucleic acid or protein compositions. An "isolated" or "purified" nucleic acid molecule or protein, or biologically active portion thereof, is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic

DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. A protein that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, 5%, (by dry weight) of contaminating protein. When the protein of the invention or biologically active portion thereof is recombinantly produced, preferably culture medium represents less than about 30%, 20%, 10%, or 5% (by dry weight) of chemical precursors or non-protein-of-interest chemicals.

Fragments and variants of the disclosed nucleotide sequences and proteins encoded thereby are also encompassed by the present invention. By "fragment" is intended a portion of the nucleotide sequence or a portion of the amino acid sequence and hence protein encoded thereby. Fragments of a nucleotide sequence may encode protein fragments that retain the biological activity of the native protein and hence their altered expression enhances resistance to pathogens. Alternatively, fragments of a nucleotide sequence that are useful as hybridization probes generally do not encode protein fragments retaining biological activity. Thus, fragments of a nucleotide sequence may range from at least about 20 nucleotides, about 50 nucleotides, about 100 nucleotides, and up to the full-length nucleotide sequence encoding the proteins of the invention.

A fragment of a *Mlo* nucleotide sequence that encodes a biologically active portion of a Mlo protein of the invention will encode at least 15, 20, 25, 30, 40, 50, 75, 100, 200, 250, 300, 350, 400, 450, 500, or 550 contiguous amino acids, or up to the total number of amino acids present in a full-length MLO protein of the invention. (For example, 509, 499, and 492, amino acids for SEQ ID NOS: 6, 12, and 14 respectively.) Fragments of a *Mlo* nucleotide sequence that are useful as hybridization probes for PCR primers generally need not encode a biologically active portion of a MLO protein.

Thus, a fragment of a *Mlo* nucleotide sequence may encode a biologically active portion of a MLO protein, or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below.

A biologically active portion of a MLO protein can be prepared by isolating a portion of one of the *Mlo* nucleotide sequences of the invention, expressing the encoded portion of the MLO protein (e.g., by recombinant expression *in vitro*), and assessing the activity of the encoded portion of the MLO protein. Nucleic acid molecules that are fragments of a *Mlo* nucleotide sequence comprise at least 16, 20, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, and 2,000 nucleotides, or up to the number of nucleotides present in a full-length *Mlo* nucleotide sequence disclosed herein.

By "variants" is intended substantially similar sequences. For nucleotide sequences, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the MLO polypeptides of the invention. Naturally occurring allelic variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis but which still encode a MLO protein of the invention. Generally, nucleotide sequence variants of the invention will have at least 40%, 50%, 60%, 70%, generally, 80%, preferably 85%, 90%, up to 95%, 98% sequence identity to its respective native nucleotide sequence.

By "variant" protein is intended a protein derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Such variants may result from, for example, genetic polymorphism or from human manipulation.

The proteins of the invention may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of the Mlo proteins can be prepared by mutations in the DNA.

Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods in Enzymol.* 154:367-382; US Patent No. 4,873,192; Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff *et al.* (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferred.

Thus, the genes and nucleotide sequences of the invention include both the naturally occurring sequences as well as mutant forms. Likewise, the proteins of the invention encompass both naturally occurring proteins as well as variations and modified forms thereof. Such variants will continue to possess the desired ability to enhance resistance to pathogens when their expression is altered. Obviously, the mutations that will be made in the DNA encoding the variant must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. See, EP Patent Application Publication No. 75,444.

The deletions, insertions, and substitutions of the protein sequences encompassed herein are not expected to produce radical changes in the characteristics of the protein. However, when it is difficult to predict the exact effect of the substitution, deletion, or insertion in advance of doing so, one skilled in the art will appreciate that the effect will be evaluated by routine screening assays. That is, the activity can be evaluated by an enhanced resistance to pathogens when the expression of the protein sequences is altered.

Variant nucleotide sequences and proteins also encompass sequences and proteins derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, one or more different *Mlo* coding sequences can be manipulated to create a new *Mlo* coding sequence possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising

sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. For example, using this approach, sequence motifs encoding a domain of interest may be shuffled between the *Mlo* gene of the invention and other known genes involved in pathogen resistance to obtain a new gene coding for a protein with an improved property of interest, such as an increased K_m in the case of an enzyme. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Cramer *et al.* (1997) *Nature Biotech.* 15:436-438; Moore *et al.* (1997) *J. Mol. Biol.* 272:336-347; Zhang *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Cramer *et al.* (1998) *Nature* 391:288-291; and U.S. Patent Nos. 5,605,793 and 5,837,458.

The maize *Mlo* homologues are associated with disease related loci as shown in Table 3. Mapping information of the sequences are given in Table 5. The maize homologue proteins are predicted to be membrane-anchored by at least six, and possibly seven membrane-spanning helices. The maize sequences find use in negative control function of the MLO protein in leaf cell death and in the onset of pathogen defense. Generally, the methods of the invention take advantage of the absence of MLO to prime responsiveness of the plant to disease.

The nucleotide sequences of the invention can be used to isolate corresponding sequences from other organisms, particularly other plants, more particularly other monocots. In this manner, methods such as PCR, hybridization, and the like can be used to identify such sequences based on their sequence homology to the sequences set forth herein. Sequences isolated based on their sequence identity to the entire *Mlo* sequences set forth herein or to fragments thereof are encompassed by the present invention.

In a PCR approach, oligonucleotide primers can be designed for use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA extracted from any plant of interest. Methods for designing PCR primers and PCR cloning are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York). See also Innis *et al.*,

eds. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, New York); Innis and Gelfand, eds. (1995) *PCR Strategies* (Academic Press, New York); and Innis and Gelfand, eds. (1999) *PCR Methods Manual* (Academic Press, New York). Known methods of PCR include, but are not
5 limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially-mismatched primers, and the like.

In hybridization techniques, all or part of a known nucleotide sequence is used as a probe that selectively hybridizes to other corresponding nucleotide
10 sequences present in a population of cloned genomic DNA fragments or cDNA fragments (*i.e.*, genomic or cDNA libraries) from a chosen organism. The hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments, or other oligonucleotides, and may be labeled with a detectable group such as ³²P, or any other detectable marker. Thus, for example, probes
15 for hybridization can be made by labeling synthetic oligonucleotides based on the *Mlo* sequences of the invention. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press,
20 Plainview, New York).

For example, the entire sequence of the *Mlo* nucleotide sequences disclosed herein, or one or more portions thereof, may be used as a probe capable of specifically hybridizing to corresponding *Mlo* sequences and messenger RNAs. To achieve specific hybridization under a variety of
25 conditions, such probes include sequences that are unique among *Mlo* sequences and are preferably at least about 10 nucleotides in length, and most preferably at least about 20 nucleotides in length. Such probes may be used to amplify corresponding *Mlo* sequences from a chosen plant by PCR. This technique may be used to isolate additional coding sequences from a desired
30 plant or as a diagnostic assay to determine the presence of coding sequences in a plant. Hybridization techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, for example, Sambrook *et al.*

(1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

Hybridization of such sequences may be carried out under stringent conditions. By "stringent conditions" or "stringent hybridization conditions" is intended conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284: $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$; where M is the

molarity of monovalent cations, %GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1°C for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with $\geq 90\%$ identity are sought, the T_m can be decreased 10°C. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4°C lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C (aqueous solution) or 32°C (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

In general, sequences that encode for a Mlo protein and hybridize to the Mlo sequences disclosed herein will be at least 40% to 50% homologous, about 60% to 70% homologous, and even about 80%, 85%, 90%, 95% to 98% homologous or more with the disclosed sequences. That is, the sequence

similarity of sequences may range, sharing at least about 40% to 50%, about 60% to 70%, and even about 80%, 85%, 90%, 95% to 98% sequence similarity.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) "reference sequence",
5 (b) "comparison window", (c) "sequence identity", (d) "percentage of sequence identity", and (e) "substantial identity".

(a) As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length
10 cDNA or gene sequence, or the complete cDNA or gene sequence.

(b) As used herein, "comparison window" makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not
15 comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically
20 introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison may be conducted by the local homology algorithm of Smith *et al.* (1981) *Adv. Appl. Math.* 2:482; by the homology alignment algorithm of Needleman *et al.* (1970) *J. Mol. Biol.*
25 48:443; by the search for similarity method of Pearson *et al.* (1988) *Proc. Natl. Acad. Sci.* 85:2444; by computerized implementations of these algorithms, including, but not limited to: CLUSTAL in the PC/Gene program by Intelligenetics, Mountain View, California; GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer
30 Group (GCG), 575 Science Drive, Madison, Wisconsin, USA; the CLUSTAL program is well described by Higgins *et al.* (1988) *Gene* 73:237-244 (1988); Higgins *et al.* (1989) *CABIOS* 5:151-153; Corpet *et al.* (1988) *Nucleic Acids Res.* 16:10881-90; Huang *et al.* (1992) *Computer Applications in the*

Biosciences 8:155-65, and Person *et al.* (1994) *Meth. Mol. Biol.* 24:307-331; preferred computer alignment methods also include the BLASTP, BLASTN, and BLASTX algorithms (see Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410). Alignment is also often performed by inspection and manual alignment.

- 5 (c) As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not
- 10 identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct
- 15 for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity.
- 20 Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).
- 25 (d) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or
- 30 deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total

number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

(e)(i) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90%, and most preferably at least 95%, compared to a reference sequence using one of the alignment programs described using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 60%, more preferably at least 70%, 80%, 90%, and most preferably at least 95%.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. However, stringent conditions encompass temperatures in the range of about 1°C to about 20°C, depending upon the desired degree of stringency as otherwise qualified herein. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides they encode are substantially identical. This may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. One indication that two nucleic acid sequences are substantially identical is when the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid.

(e)(ii) The term "substantial identity" in the context of a peptide indicates that a peptide comprises a sequence with at least 70% sequence identity to a reference sequence, preferably 80%, more preferably 85%, most preferably at least 90% or 95% sequence identity to the reference sequence over a specified comparison window. Preferably, optimal alignment is conducted using the homology alignment algorithm of Needleman *et al.* (1970) *J. Mol.*

Biol. 48:443. An indication that two peptide sequences are substantially identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. Thus, a peptide is substantially identical to a second peptide, for example, where the two peptides differ only by a conservative substitution. Peptides that are "substantially similar" share sequences as noted above except that residue positions that are not identical may differ by conservative amino acid changes.

The invention is drawn to methods for creating or enhancing resistance in a plant to plant pests by modulating the activity of *Mlo* genes in the plant. While the invention is not bound by any particular mechanism of action, it is believed that the methods of the invention will result in broad-based resistance in the modified plant. Accordingly, the methods are also useful in protecting plants against fungal pathogens, viruses, nematodes, insects and the like.

By "modulating activity" is intended that the expression of the *Mlo* gene is altered in some manner. Such modulation (increase or decrease) of expression results in enhanced resistance to pathogens. Generally, the methods of the invention will result in a decrease in the native protein or in protein activity. Thus, plants and plant cells are obtained having altered levels of MLO protein, preferably a decrease in protein levels. Such plants, plant cells and plant tissues are "modified" in that MLO protein levels are altered. As noted below, various methods are available for creating modified plants, plant cells and plant tissues including transformation and transfection leading to altered *Mlo* expression in the modified plant, plant cell or tissue.

The invention is drawn to compositions and methods for inducing resistance in a plant to plant pests. Accordingly, the compositions and methods are also useful in protecting plants against fungal pathogens, viruses, nematodes, insects and the like.

By "disease resistance" is intended that the plants avoid the disease symptoms that are the outcome of plant-pathogen interactions. That is, pathogens are prevented from causing plant diseases and the associated disease symptoms, or alternatively, the disease symptoms caused by the pathogen is minimized or lessened. The methods of the invention can be utilized to protect

plants from disease, particularly those diseases that are caused by plant pathogens.

Pathogens of the invention include, but are not limited to, viruses or viroids, bacteria, insects, nematodes, fungi, and the like. Viruses include any plant virus, for example, tobacco or cucumber mosaic virus, ringspot virus, necrosis virus, maize dwarf mosaic virus, etc. Specific fungal and viral pathogens for the major crops include: Soybeans: *Phytophthora megasperma* fsp. *glycinea*, *Macrophomina phaseolina*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Fusarium oxysporum*, *Diaporthe phaseolorum* var. *sojae* (10 *Phomopsis sojae*), *Diaporthe phaseolorum* var. *caulivora*, *Sclerotium rolfsii*, *Cercospora kikuchii*, *Cercospora sojae*, *Peronospora manshurica*, *Colletotrichum dematium* (*Colletotrichum truncatum*), *Corynespora cassicola*, *Septoria glycines*, *Phyllosticta sojicola*, *Alternaria alternata*, *Pseudomonas syringae* p.v. *glycinea*, *Xanthomonas campestris* p.v. *phaseoli*, *Microsphaera* (15 *diffusa*, *Fusarium semitectum*, *Phialophora gregata*, Soybean mosaic virus, *Glomerella glycines*, Tobacco Ring spot virus, Tobacco Streak virus, *Phakopsora pachyrhizi*, *Pythium aphanidermatum*, *Pythium ultimum*, *Pythium debaryanum*, Tomato spotted wilt virus, *Heterodera glycines* *Fusarium solani*; Canola: *Albugo candida*, *Alternaria brassicae*, *Leptosphaeria maculans*, (20 *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Mycosphaerella brassicicola*, *Pythium ultimum*, *Peronospora parasitica*, *Fusarium roseum*, *Alternaria alternata*; Alfalfa: *Clavibacter michiganense* subsp. *insidiosum*, *Pythium ultimum*, *Pythium irregulare*, *Pythium splendens*, *Pythium debaryanum*, *Pythium aphanidermatum*, *Phytophthora megasperma*, *Peronospora* (25 *trifoliorum*, *Phoma medicaginis* var. *medicaginis*, *Cercospora medicaginis*, *Pseudopeziza medicaginis*, *Leptotrochila medicaginis*, *Fusarium*, *Xanthomonas campestris* p.v. *alfalfae*, *Aphanomyces euteiches*, *Stemphylium herbarum*, *Stemphylium alfalfae*; Wheat: *Pseudomonas syringae* p.v. *atrofaciens*, *Urocystis agropyri*, *Xanthomonas campestris* p.v. *translucens*, *Pseudomonas* (30 *syringae* p.v. *syringae*, *Alternaria alternata*, *Cladosporium herbarum*, *Fusarium graminearum*, *Fusarium avenaceum*, *Fusarium culmorum*, *Ustilago tritici*, *Ascochyta tritici*, *Cephalosporium gramineum*, *Collotetrichum graminicola*, *Erysiphe graminis* f.sp. *tritici*, *Puccinia graminis* f.sp. *tritici*,

- Puccinia recondita* f.sp. *tritici*, *Puccinia striiformis*, *Pyrenophora tritici-repentis*, *Septoria nodorum*, *Septoria tritici*, *Septoria avenae*,
Pseudocercospora herpotrichoides, *Rhizoctonia solani*, *Rhizoctonia cerealis*,
Gaeumannomyces graminis var. *tritici*, *Pythium aphanidermatum*, *Pythium*
5 *arrhenomanes*, *Pythium ultimum*, *Bipolaris sorokiniana*, Barley Yellow Dwarf
Virus, Brome Mosaic Virus, Soil Borne Wheat Mosaic Virus, Wheat Streak
Mosaic Virus, Wheat Spindle Streak Virus, American Wheat Striate Virus,
Claviceps purpurea, *Tilletia tritici*, *Tilletia laevis*, *Ustilago tritici*, *Tilletia*
indica, *Rhizoctonia solani*, *Pythium arrhenomannes*, *Pythium graminicola*,
10 *Pythium aphanidermatum*, High Plains Virus, European wheat striate virus;
Sunflower: *Plasmophora halstedii*, *Sclerotinia sclerotiorum*, Aster Yellows,
Septoria helianthi, *Phomopsis helianthi*, *Alternaria helianthi*, *Alternaria*
zinniae, *Botrytis cinerea*, *Phoma macdonaldii*, *Macrophomina phaseolina*,
Erysiphe cichoracearum, *Rhizopus oryzae*, *Rhizopus arrhizus*, *Rhizopus*
15 *stolonifer*, *Puccinia helianthi*, *Verticillium dahliae*, *Erwinia carotovorum* pv.
carotovora, *Cephalosporium acremonium*, *Phytophthora cryptogea*, *Albugo*
tragopogonis; Corn: *Fusarium moniliforme* var. *subglutinans*, *Erwinia*
stewartii, *Fusarium moniliforme*, *Gibberella zeae* (*Fusarium graminearum*),
Stenocarpella maydi (*Diplodia maydis*), *Pythium irregulare*, *Pythium*
20 *debaryanum*, *Pythium graminicola*, *Pythium splendens*, *Pythium ultimum*,
Pythium aphanidermatum, *Aspergillus flavus*, *Bipolaris maydis* O, T
(*Cochliobolus heterostrophus*), *Helminthosporium carbonum* I, II & III
(*Cochliobolus carbonum*), *Exserohilum turcicum* I, II & III, *Helminthosporium*
pedicellatum, *Physoderma maydis*, *Phyllosticta maydis*, *Kabatiella-maydis*,
25 *Cercospora sorghi*, *Ustilago maydis*, *Puccinia sorghi*, *Puccinia polysora*,
Macrophomina phaseolina, *Penicillium oxalicum*, *Nigrospora oryzae*,
Cladosporium herbarum, *Curvularia lunata*, *Curvularia inaequalis*, *Curvularia*
pallens, *Clavibacter michiganense* subsp. *nebraskense*, *Trichoderma viride*,
Maize Dwarf Mosaic Virus A & B, Wheat Streak Mosaic Virus, Maize
30 Chlorotic Dwarf Virus, *Claviceps sorghi*, *Pseudonomas avenae*, *Erwinia*
chrysanthemi pv. *zea*, *Erwinia carotovora*, *Corn stunt spiroplasma*, *Diplodia*
macrospora, *Sclerophthora macrospora*, *Peronosclerospora sorghi*,
Peronosclerospora philippinensis, *Peronosclerospora maydis*,

- Peronosclerospora sacchari*, *Sphacelotheca reiliana*, *Physopella zae*,
Cephalosporium maydis, *Cephalosporium acremonium*, Maize Chlorotic Mottle
 Virus, High Plains Virus, Maize Mosaic Virus, Maize Rayado Fino Virus,
 Maize Streak Virus, Maize Stripe Virus, Maize Rough Dwarf Virus; Sorghum:
- 5 *Exserohilum turcicum*, *Colletotrichum graminicola* (*Glomerella graminicola*),
Cercospora sorghi, *Gloeocercospora sorghi*, *Ascochyta sorghina*,
Pseudomonas syringae p.v. *syringae*, *Xanthomonas campestris* p.v. *holcicola*,
Pseudomonas andropogonis, *Puccinia purpurea*, *Macrophomina phaseolina*,
Perconia circinata, *Fusarium moniliforme*, *Alternaria alternata*, *Bipolaris*
 10 *sorghicola*, *Helminthosporium sorghicola*, *Curvularia lunata*, *Phoma insidiosa*,
Pseudomonas avenae (*Pseudomonas alboprecipitans*), *Ramulispora sorghi*,
Ramulispora sorghicola, *Phyllachara sacchari*, *Sporisorium reilianum*
 (*Sphacelotheca reiliana*), *Sphacelotheca cruenta*, *Sporisorium sorghi*,
 Sugarcane mosaic H, Maize Dwarf Mosaic Virus A & B, *Claviceps sorghi*,
 15 *Rhizoctonia solani*, *Acremonium strictum*, *Sclerophthora macrospora*,
Peronosclerospora sorghi, *Peronosclerospora philippinensis*, *Sclerospora*
graminicola, *Fusarium graminearum*, *Fusarium oxysporum*, *Pythium*
arrhenomanes, *Pythium graminicola*, etc.

- Nematodes include parasitic nematodes such as root-knot, cyst, lesion,
 20 and renniform nematodes, etc.

- Insect pests include insects selected from the orders Coleoptera, Diptera,
 Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera,
 Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc.,
 particularly Coleoptera and Lepidoptera. Insect pests of the invention for the
 25 major crops include: Maize: *Ostrinia nubilalis*, European corn borer; *Agrotis*
ipsilon, black cutworm; *Helicoverpa zea*, corn earworm; *Spodoptera*
frugiperda, fall armyworm; *Diatraea grandiosella*, southwestern corn borer;
Elasmopalpus lignosellus, lesser cornstalk borer; *Diatraea saccharalis*,
 sugarcane borer; *Diabrotica virgifera*, western corn rootworm; *Diabrotica*
 30 *longicornis barberi*, northern corn rootworm; *Diabrotica undecimpunctata*
howardi, southern corn rootworm; *Melanotus spp.*, wireworms; *Cyclocephala*
borealis, northern masked chafer (white grub); *Cyclocephala immaculata*,
 southern masked chafer (white grub); *Popillia japonica*, Japanese beetle;

- Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Anuraphis maidiradicis*, corn root aphid; *Blissus leucopterus leucopterus*, chinch bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus sanguinipes*, migratory grasshopper;
- 5 *Hylemya platura*, seedcorn maggot; *Agromyza parvicornis*, corn blot leafminer; *Anaphothrips obscurus*, grass thrips; *Solenopsis milesta*, thief ant; *Tetranychus urticae*, twospotted spider mite; Sorghum: *Chilo partellus*, sorghum borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Feltia subterranea*, granulate
- 10 cutworm; *Phyllophaga crinita*, white grub; *Eleodes*, *Conoderus*, and *Aeolus* spp., wireworms; *Oulema melanopus*, cereal leaf beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid; *Sipha flava*, yellow sugarcane aphid; *Blissus leucopterus leucopterus*, chinch bug; *Contarinia sorghicola*, sorghum
- 15 midge; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Wheat: *Pseudaletia unipunctata*, army worm; *Spodoptera frugiperda*, fall armyworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Agrotis orthogonia*, western cutworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Oulema melanopus*, cereal leaf beetle;
- 20 *Hypera punctata*, clover leaf weevil; *Diabrotica undecimpunctata howardi*, southern corn rootworm; Russian wheat aphid; *Schizaphis graminum*, greenbug; *Macrosiphum avenae*, English grain aphid; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Melanoplus sanguinipes*, migratory grasshopper; *Mayetiola destructor*, Hessian
- 25 fly; *Sitodiplosis mosellana*, wheat midge; *Meromyza americana*, wheat stem maggot; *Hylemya coarctata*, wheat bulb fly; *Frankliniella fusca*, tobacco thrips; *Cephus cinctus*, wheat stem sawfly; *Aceria tulipae*, wheat curl mite; Sunflower: *Suleima helianthana*, sunflower bud moth; *Homoeosoma electellum*, sunflower moth; *zygogramma exclamationis*, sunflower beetle; *Bothyrus gibbosus*, carrot
- 30 beetle; *Neolasioptera murtfeldtiana*, sunflower seed midge; Cotton: *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Spodoptera exigua*, beet armyworm; *Pectinophora gossypiella*, pink bollworm; *Anthonomus grandis grandis*, boll weevil; *Aphis gossypii*, cotton aphid; *Pseudatomoscelis*

default cell suicide program in foliar tissue. Also, the MLO protein could have a specific negative regulatory function which works by down-regulating multiple disease-related functions. In this instance, spontaneous cell death in *mlo* mutant genotypes merely represents cell death because of accumulating
5 activation of defense responses.

Several methods are available in the art for modulating the activity of *Mlo* genes. *Mlo* antisense sequences can be expressed in the plant cell. Such sequences will function to decrease expression of the maize *Mlo* genes as well as *Mlo* genes in other plants where the *Mlo* sequences share sequence identity.

10 It is recognized that with these nucleotide sequences, antisense constructions, complementary to at least a portion of the messenger RNA (mRNA) for the *Mlo* sequences can be constructed. Antisense nucleotides are constructed to hybridize with the corresponding mRNA. Modifications of the antisense sequences may be made as long as the sequences hybridize to and
15 interfere with expression of the corresponding mRNA. In this manner, antisense constructions having 70%, preferably 80%, more preferably 85% sequence similarity to the corresponding antisensed sequences may be used. Furthermore, portions of the antisense nucleotides may be used to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleotides,
20 100 nucleotides, 200 nucleotides, or greater may be used.

The antisense construct may be proximal to the 5'-terminus or capping site, downstream from the capping site, between the capping site and the initiation codon and may cover all or a portion of the non-coding region. Additionally, the sequence may bridge the non-coding and coding region, be
25 complementary to all or part of the coding region, to the 3'-terminus of the coding region, or complementary to the 3'-untranslated region of the mRNA.

It is recognized that the particular site to which the antisense sequence binds and the length of the antisense sequence will vary depending upon the degree of inhibition desired, the uniqueness of the sequence, the stability of the
30 antisense sequence, and the like. See, for example, U.S. Patent Nos. 5,453,566; 5,530,192; and 5,728,926; all of which are herein incorporated by reference.

The nucleotide sequences of the present invention may also be used in the sense orientation to suppress the expression of endogenous genes in plants.

seriatus, cotton fleahopper; *Trialeurodes abutilonea*, bandedwinged whitefly;
Lygus lineolaris, tarnished plant bug; *Melanoplus femurrubrum*, redlegged
 grasshopper; *Melanoplus differentialis*, differential grasshopper; *Thrips tabaci*,
 onion thrips; *Frankliniella fusca*, tobacco thrips; *Tetranychus cinnabarinus*,
 5 carmine spider mite; *Tetranychus urticae*, twospotted spider mite; Rice:
Diatraea saccharalis, sugarcane borer; *Spodoptera frugiperda*, fall armyworm;
Helicoverpa zea, corn earworm; *Colaspis brunnea*, grape colaspis;
Lissorhoptrus oryzophilus, rice water weevil; *Sitophilus oryzae*, rice weevil;
Nephotettix nigropictus, rice leafhopper; *Blissus leucopterus leucopterus*,
 10 chinch bug; *Acrosternum hilare*, green stink bug; Soybean: *Pseudoplusia*
inclusens, soybean looper; *Anticarsia gemmatilis*, velvetbean caterpillar;
Plathypena scabra, green cloverworm; *Ostrinia nubilalis*, European corn borer;
Agrotis ipsilon, black cutworm; *Spodoptera exigua*, beet armyworm; *Heliothis*
virescens, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Epilachna*
 15 *varivestis*, Mexican bean beetle; *Myzus persicae*, green peach aphid; *Empoasca*
fabae, potato leafhopper; *Acrosternum hilare*, green stink bug; *Melanoplus*
femurrubrum, redlegged grasshopper; *Melanoplus differentialis*, differential
 grasshopper; *Hylemya platura*, seedcorn maggot; *Sericothrips variabilis*,
 soybean thrips; *Thrips tabaci*, onion thrips; *Tetranychus turkestanii*, strawberry
 20 spider mite; *Tetranychus urticae*, twospotted spider mite; Barley: *Ostrinia*
nubilalis, European corn borer; *Agrotis ipsilon*, black cutworm; *Schizaphis*
graminum, greenbug; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum*
hilare, green stink bug; *Euschistus servus*, brown stink bug; *Delia platura*,
 seedcorn maggot; *Mayetiola destructor*, Hessian fly; *Petrobia latens*, brown
 25 wheat mite; Oil Seed Rape: *Brevicoryne brassicae*, cabbage aphid; *Phyllotreta*
cruciferae, Flea beetle; *Mamestra configurata*, Bertha armyworm; *Plutella*
xylostella, Diamond-back moth; *Delia* ssp., Root maggots.

The present invention exploits the use of *Mlo* genes. The *Mlo* sequences
 provided herein can be utilized to alter the expression of native *Mlo* genes in
 30 plants. The *mlo* mutation confers recessive resistance to pathogens. Broad
 spectrum resistance in plants can be enhanced by defective *Mlo* genes.

While the invention is not bound by any model, *Mlo* could have a
 negative control function in leaf cell death. In this model, *Mlo* would suppress a

Methods for suppressing gene expression in plants using nucleotide sequences in the sense orientation are known in the art. The methods generally involve transforming plants with a DNA construct comprising a promoter that drives expression in a plant operably linked to at least a portion of a nucleotide
5 sequence that corresponds to the transcript of the endogenous gene. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, preferably greater than about 65% sequence identity, more preferably greater than about 85% sequence identity, most preferably greater than about 95% sequence identity. See, U.S. Patent
10 Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

Individual *Mlo* genes or sets of *Mlo* genes can be rendered nonfunctional by mutagenesis. Such mutagenesis techniques include transposon disruption and recovery of such disruptions by reverse genetics approaches. Likewise, transformation-mediated mutagenesis may be utilized.

15 Disruption may be accomplished by transformation with gene replacement or gene truncation-disruption. See, for example, Bowen *et al.* (1995) *Mol. Gen. Gent.* 246:196-205; Walz *et al.* (1993) *Curr. Gent.* 25:421-427; Sweigard *et al.* (1992) *Mol. Gen. Gent.* 232:183-190; Hohn and Desjardins (1992) *Mol. Plant Microbe Interact* 5:249-256; Weber and Laitner (1994) *Curr.*
20 *Gent.* 26:461-467; Templeton *et al.* (1994) *Gene* 142:141-146; Gorlach *et al.* (1998) *Appl. Environ. Microbiol.* 64:385-391; Schaeffer *et al.* (1994) *Appl. Environ. Microbiol.* 60:594-598; and the like, herein incorporated by reference. See also, Kempin *et al.* (1997) *Nature* 389:802-803 and Koncz *et al.* (1992) *Plant Mol. Biol.* 20:963-976.

25 Alternatively, the naturally occurring *Mlo* sequence or sequences may be modified by site-directed mutagenesis. Such methods may be utilized to induce specific alterations in targeted genes. One means for site-directed mutagenesis includes targeting modification or mutation of the *Mlo* sequences by homologous recombination. The method involves the use of RNA-DNA
30 hybrid oligonucleotides. Such nucleotides exploit the natural recombinogenicity of RNA-DNA hybrids. The oligonucleotides are duplex oligonucleotides that share homology with at least one *Mlo* sequence. While any region of the *Mlo* sequence can be targeted, it may be preferable to target

the 5' region of the *Mlo* sequence. See, for example, U.S. Patent No. 5,565,350 that describes chimeric oligonucleotides useful for targeted gene correction for use in cultured mammalian cells; as well as U.S. provisional application Serial No. 06/065,628 drawn to gene manipulation in plant cells, herein incorporated
5 by reference. Such methods can be used to alter or disrupt the ATG start codon for the gene.

Alternatively, protein coding regions of *Mlo* genes can be altered in such a manner that the gene products or proteins perform their function in a dominant negative manner resulting in a resistant phenotype. See, for example Krylov *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:12274-12279.
10

Where the method of the invention relies upon the expression of an altered *Mlo* sequence or an *Mlo* antisense sequence in a plant, a number of promoters can be used. The promoters can be selected from constitutive and/or inducible promoters. Such promoters include those from pathogenesis-related
15 proteins (PR proteins), which are induced following infection by a pathogen; e.g., PR proteins, SAR proteins, beta-1,3-glucanase, chitinase, etc. See, for example, Redolfi *et al.* (1983) *Neth. J. Plant Pathol.* 89:245-254; Uknes *et al.* (1992) *The Plant Cell* 4:645-656; and Van Loon (1985) *Plant Mol. Virol.* 4:111-116. See also copending application entitled "Inducible Maize promoters",
20 filed and herein incorporated by reference.

Of interest are promoters that are expressed locally at or near the site of pathogen infection. See, for example, Marineau *et al.* (1987) *Plant Mol. Biol.* 9:335-342; Matton *et al.* (1989) *Molecular Plant-Microbe Interactions* 2:325-331; Somsisch *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:2427-2430;
25 Somsisch *et al.* (1988) *Molecular and General Genetics* 2:93-98; and Yang, Y (1996) *Proc. Natl. Acad. Sci. USA* 93:14972-14977. See also, Chen *et al.* (1996) *Plant J.* 10:955-966; Zhang and Sing (1994) *Proc. Natl. Acad. Sci. USA* 91:2507-2511; Warner *et al.* (1993) *Plant J.* 3:191-201; Siebertz *et al.* (1989) *Plant Cell* 1:961-968; and the references cited therein. Of particular interest is
30 the inducible promoter for the maize PRms gene, whose expression is induced by the pathogen *Fusarium moniliforme* (see, for example, Cordero *et al.* (1992) *Physiological and Molecular Plant Pathology* 41:189-200).

Constitutive promoters include, for example, the core promoter of the Rsyn7 (copending application serial number 08/661,601), the 35S promoter, the core 35S promoter, and the like. Other constitutive promoters include, for example, U.S. Patent Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; 5,608,142. See also, copending application
5 5,466,785; 5,399,680; 5,268,463; 5,608,142. See also, copending application entitled "Constitutive Maize Promoters" Provisional Application Serial No. 60/076,075 filed February 26, 1998, and herein incorporated by reference.

Tissue specific promoters include Yamamoto *et al.* (1997) *Plant J.* 12(2):255-265; Kawamata *et al.* (1997) *Plant Cell Physiol.* 38(7):792-803;
10 Hansen *et al.* (1997) *Mol. Gen Genet.* 254(3):337-343; Russell *et al.* (1997) *Transgenic Res.* 6(2):157-168; Rinehart *et al.* (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp *et al.* (1996) *Plant Physiol.* 112(2):525-535; Canevascini *et al.* (1996) *Plant Physiol.* 112(2):513-524; Yamamoto *et al.* (1994) *Plant Cell Physiol.* 35(5):773-778; Lam (1994) *Results Probl. Cell*
15 *Differ.* 20:181-196; Orozco *et al.* (1993) *Plant Mol. Biol.* 23(6):1129-1138; Matsuoka *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90(20):9586-9590; and Guevara-Garcia *et al.* (1993) *Plant J.* 4(3):495-505.

The methods of the invention can be used with other methods for increasing pathogen resistance in plants. See, for example, Cai *et al.* (1997)
20 *Science* 275:832-834; Roberts and Galloway (1984) *J. Heredity* 75:147-148; Ryerson and Heath (1996) *Plant Cell* 8:393-402 and Dangl *et al.* (1996) *Plant Cell* 8:1793-1807.

Altered *Mlo* sequences or antisense *Mlo* sequences of the invention can be introduced into any plant. The sequences to be introduced may be used in
25 expression cassettes for expression in any plant of interest where expression in the plant cell is necessary. In other instances, such as for recombination, oligonucleotides are synthesized, purified and introduced into the plant cell.

Where expression cassettes are needed, such expression cassettes will comprise a transcriptional initiation region linked to the coding sequence or
30 antisense sequence of the nucleotide of interest. Such an expression cassette is provided with a plurality of restriction sites for insertion of the sequence to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

The transcriptional initiation region, the promoter, may be native or analogous or foreign or heterologous to the plant host. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. By foreign is intended that the transcriptional initiation region is not found in the native plant into which the transcriptional initiation region is introduced. As used herein a chimeric gene comprises a coding sequence operably linked to a transcription initiation region that is heterologous to the coding sequence.

While it may be preferable to express the sequences using heterologous promoters, the native promoter sequences may be used. Such constructs would change expression levels of a MLO protein in the plant or plant cell. Thus, the phenotype of the plant or plant cell is altered.

The transcriptional cassette will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region, a DNA sequence of interest, and a transcriptional and translational termination region functional in plants. The termination region may be native with the transcriptional initiation region, may be native with the DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell.* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158; Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; Joshi *et al.* (1987) *Nucleic Acid Res.* 15:9627-9639.

Nucleotide sequences of the invention are provided in expression cassettes for expression in the plant of interest. The cassette will include 5' and 3' regulatory sequences operably linked to the sequence of interest. The cassette may additionally contain at least one additional sequence to be cotransformed into the organism. Alternatively, the additional sequence(s) can be provided on another expression cassette.

Where appropriate, the gene(s) may be optimized for increased expression in the transformed plant. That is, the genes can be synthesized using plant-preferred codons for improved expression. See, for example, Campbell

and Gowri (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, 5,436,391, and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

5 Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene expression. The G-C content of the sequence may be adjusted to levels
10 average for a given cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance
15 translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein *et al.* (1989) *PNAS USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986); MDMV leader (Maize Dwarf Mosaic Virus); *Virology* 154:9-20), and human
20 immunoglobulin heavy-chain binding protein (BiP), (Macejak *et al.* (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling *et al.* (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie *et al.* (1989) in *Molecular Biology of RNA*, ed. Cech (Liss, New York), pp. 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also, Della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968. Other methods known to
25 enhance translation can also be utilized, for example, introns, and the like.

In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation
30 and, as appropriate, in the proper reading frame. Towards this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in*

vitro mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g. transitions and transversions, may be involved.

The sequences of the present invention can be used to transform or transfect any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained. Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation (Townsend *et al.*, U.S. Pat No. 5,563,055), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.*, U.S. Patent No. 4,945,050; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Ann. Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize); Tomes, U.S. Patent No. 5,240,855; Buising *et al.*, U.S. Patent Nos. 5,322,783 and 5,324,646; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg (Springer-Verlag, Berlin) (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren *et al.* (1984) *Nature (London)* 311:763-764; Bowen *et al.*, U.S. Patent No. 5,736,369 (cereals); Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA*

84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418 and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

10 The modified plant may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports*, 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having the desired phenotypic characteristic identified. Two or more generations may
15 be grown to ensure that the subject phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure the desired phenotype or other property has been achieved.

 The *Mlo* sequences of the invention are also useful as molecular markers. Such markers are useful in breeding programs, particularly those
20 aimed at improving disease resistance. The maize *Mlo* sequences have been mapped to chromosome locations and these positions related to known disease resistance loci.

 The present invention provides a method of genotyping a plant comprising a polynucleotide of the present invention. Preferably, the plant is a
25 monocot, such as maize or sorghum. Genotyping provides a means of distinguishing homologs of a chromosome pair and can be used to differentiate segregants in a plant population. Molecular marker methods can be used for phylogenetic studies, characterizing genetic relationships among crop varieties, identifying crosses or somatic hybrids, localizing chromosomal segments
30 affecting monogenic traits, map based cloning, and the study of quantitative inheritance. See, e.g., *Plant Molecular Biology: A Laboratory Manual*, Chapter 7, Clark, Ed., Springer-Verlag, Berlin (1997). For molecular marker methods, see generally, *The DNA Revolution* by Andrew H. Paterson (1996)

(Chapter 2) in: *Genome Mapping in Plants* (ed. Andrew H. Paterson) by Academic Press/R. G. Landis Company, Austin, Texas, pp. 7-21.

The particular method of genotyping in the present invention may employ any number of molecular marker analytic techniques such as, but not limited to, restriction fragment length polymorphisms (RFLPs). RFLPs are the product of allelic differences between DNA restriction fragments caused by nucleotide sequence variability. As is well known to those of skill in the art, RFLPs are typically detected by extraction of genomic DNA and digestion with a restriction enzyme. Generally, the resulting fragments are separated according to size and hybridized with a probe; single copy probes are preferred. Restriction fragments from homologous chromosomes are revealed. Differences in fragment size among alleles represent an RFLP. Thus, the present invention further provides a means to follow segregation of a gene or nucleic acid of the present invention as well as chromosomal sequences genetically linked to these genes or nucleic acids using such techniques as RFLP analysis. Linked chromosomal sequences are within 50 centiMorgans (Cm), often within 40 or 30 Cm, preferably within 20 or 10 Cm, more preferably within 5, 3, 2, or 1 Cm of a gene.

In the present invention, the nucleic acid probes employed for molecular marker mapping of plant nuclear genomes selectively hybridize, under selective hybridization conditions, to a gene encoding a polynucleotide of the present invention. In preferred embodiments, the probes are selected from polynucleotides of the present invention. Typically, these probes are cDNA probes or *PstI* genomic clones. The length of the probes is discussed in greater detail, *supra*, but are typically at least 15 bases in length, more preferably at least 20, 25, 30, 35, 40, or 50 bases in length. Generally, however, the probes are less than about 1 kilobase in length. Preferably, the probes are single copy probes that hybridize to a unique locus in a haploid chromosome complement. Some exemplary restriction enzymes employed in RFLP mapping are *EcoRI*, *EcoRV*, and *SstI*. As used herein the term "restriction enzyme" includes reference to a composition that recognizes and, alone or in conjunction with another composition, cleaves at a specific nucleotide sequence.

The method of detecting an RFLP comprises the steps of (a) digesting genomic DNA of a plant with a restriction enzyme; (b) hybridizing a nucleic acid probe, under selective hybridization conditions, to a sequence of a polynucleotide of the present of said genomic DNA; (c) detecting therefrom a RFLP. Other methods of differentiating polymorphic (allelic) variants of polynucleotides of the present invention can be had by utilizing molecular marker techniques well known to those of skill in the art including such techniques as: 1) single-stranded conformation analysis (SSCP); 2) denaturing gradient gel electrophoresis (DGGE); 3) Rnase protection assays; 4) allele-specific oligonucleotides (ASOs); 5) the use of proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein; and 6) allele-specific PCR. Other approaches based on the detection of mismatches between the two complementary DNA strands include clamped denaturing gel electrophoresis (CDGE); heteroduplex analysis (HA); and chemical mismatch cleavage (CMC). Exemplary polymorphic variants are provided in Table I, *supra*. Thus, the present invention further provides a method of genotyping comprising the steps of contacting, under stringent hybridization conditions, a sample suspected of comprising a polynucleotide of the present invention with a nucleic acid probe. Generally, the sample is a plant sample; preferably, a sample suspected of comprising a maize polynucleotide of the present invention (e.g., gene, mRNA). The nucleic acid probe selectively hybridizes, under stringent conditions, to a subsequence of a polynucleotide of the present invention comprising a polymorphic marker. Selective hybridization of the nucleic acid probe to the polymorphic marker nucleic acid sequence yields a hybridization complex. Detection of the hybridization complex indicates the presence of that polymorphic marker in the sample. In preferred embodiments, the nucleic acid probe comprises a polynucleotide of the present invention.

The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

Mlo Sequences

Eight maize *Mlo* homologue sequences have been identified. The nucleotide sequences (SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15) and amino acid sequences (SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, and 16) are provided. DNA sequence analysis was performed according to Sambrook *et al.* (1989) *Mol. Cloning: A Laboratory Manual*, 2nd ed. (Cold Spring Harbor Laboratory Press, Plainview, New York). The maize *Mlo* homologue/orthologue cDNA sequences were identified from mRNA isolated from various maize tissues and used to make cDNAs that are then cloned into vectors, usually pSPORT1. These cDNA libraries are EST sequenced via standard dye-fluorescence labeling and ABI machine electrophoresis and image capture. The *Mlo* homologues were identified by their blast score identity to the barley HvMLO1 published sequence. These maize clones, in particular the longest members of each contig or gene, were obtained and additional sequencing was done on them using oligonucleotide primers designed to internal portions of the cDNA and dye-fluorescence labeling and ABI machine electrophoresis and image capture. The complete edited sequences were assembled and analyzed.

The sequences from maize show sequence similarity to the published barley *Mlo* sequence as provided in Tables 1 and 2. Figure 1 provides a sequence alignment for the maize *Mlo* homologues with the barley *Mlo* sequence designated as HvMlo1. Figure 2 groups the *Mlo* sequences based on relatedness as well as the plant tissue from which it was isolated.

The ZmMLO genes were mapped by RFLP analysis using Southern blots of genomic DNA isolated from F2 and F3 and F4 segregating maize populations. The DNA was isolated using a modified CTAB adapted from the CERES RFLP Lab Manuel based on the Saghai-Mahoof procedure (Saghai-Mahoof *et al.* (1984) *Proc. Natl. Acad. Sci. USA* 81:8014-8018). Genomic DNA was cut with four restriction enzymes *Bam*HI, *Eco*RI, *Eco*RV, and *Hind*III and run (5µg/lane) on 0.8% TAE agarose gels for 800 volt-hours. The gels were Southern blot transferred to Hybond-N membrane (Amersham Life Science), baked and UV-crosslinked. The ZmMLO cDNA inserts were liberated from its vector by cleaving with two restriction enzymes, usually *Sal*I

and *NotI*, and the insert was purified from agarose gels following electrophoresis. The clone was random prime labeled with ^{32}P -dCTP using the RTS (Gibco BRL) labeling kit. The cDNA inserts were used to probe parental screening blots to identify a mapping population with a RFLP polymorphism.

- 5 Once a polymorphism was identified, the inserts were used to probe mapping blots containing DNA from segregating individuals. The map position was determined using MAPMAKER/EXP 3.0 (Lander *et al* (1987) *Genomics* 1:174-181) by scoring 86 segregating progeny as homozygous parent A, as homozygous parent B, or as heterozygous. The map position was assigned to
- 10 an existing core RFLP map of either of the following three populations: ALEB9 (240 individuals) DRAG2 (283 individuals) or MARS (1075 individuals). Table 3 shows the association of maize *Mlo* homologues and disease related loci and QTLS.

TABLE I

Maize <i>MLO</i> Gene		BestFit (Similarity/Identity)								
		HvMLO1	ZmMLO1	ZmMLO2	ZmMLO4	ZmMLO5	ZmMLO6	ZmMLO7	ZmMLO8	ZmMLO9
Protein (Similarity/Identity) to:										
HvMLO1	100%									
ZmMLO1	72	67		53	35	45	58	40	44	37
ZmMLO2	61	100%		45	35	38	51	39	43	39
ZmMLO4	47	56	100%		40	48	58	43	46	40
ZmMLO5	54	44	49		100%	46	38	39	40	46
ZmMLO6	66	47	56	56	56	100%	45	50	76	41
ZmMLO7	48	61	66	66	48	51	100%	41	43	39
ZmMLO8	54	49	52	52	48	58	52	100%	56	41
ZmMLO9	48	55	56	56	52	81	54	64	100%	43
		48	49	49	51	56	50	51	56	100%

TABLE 2

Gap (Similarity/Identity)		Protein Homology (Similarity/Identity) to:								
Maize <i>MLO</i> Gene	Hv <i>MLO1</i>	Zm <i>MLO1</i>	Zm <i>MLO2</i>	Zm <i>MLO4</i>	Zm <i>MLO5</i>	Zm <i>MLO6</i>	Zm <i>MLO7</i>	Zm <i>MLO8</i>	Zm <i>MLO9</i>	
Hv <i>MLO1</i>	100%	66	43	31	33	55	36	40	32	
Zm <i>MLO1</i>	71	100%	36	24	32	43	28	37	28	
Zm <i>MLO2</i>	53	46	100%	36	36	57	39	45	34	
Zm <i>MLO4</i>	42	31	46	100%	32	34	35	37	42	
Zm <i>MLO5</i>	40	38	42	38	100%	43	42	60	36	
Zm <i>MLO6</i>	63	51	64	43	49	100%	40	41	36	
Zm <i>MLO7</i>	43	35	47	43	48	51	100%	51	40	
Zm <i>MLO8</i>	49	48	54	49	67	52	59	100%	40	
Zm <i>MLO9</i>	40	36	42	53	48	46	44	51	100%	

TABLE 3
Association of Maize MLO Homologues and Disease Related Loci and QTLs

<u>Maize MLO Homologue</u>	<u>Nearby Disease Resistance Loci or QTLs</u>
ZmMLO1 1.01	1.01 European Corn Borer QTL
	1.01/2 Northern Corn Leaf Blight QTL
ZmMLO2 1.04	1.03/6 Northern Corn Leaf Blight QTL
	1.04 Gray Leaf Spot QTL
	1.04 Maize Streak Virus (<i>msv1</i>)
	1.05 Stewart's Wilt QTL
ZmMLO4 3.05	3.04/5 European Corn Borer QTL
	3.04/5 Gibberella Stalk Rot QTL
ZmMLO5 3.06	3.07/8 Northern Corn Leaf Blight QTL
ZmMLO6a 5.05	5.04 Gibberella Stalk Rot QTL
	5.06 Northern Corn Leaf Blight QTL
ZmMLO6b 6.02/3	6.01 Southern Corn Leaf Blight (<i>rhml</i>)
ZmMLO7 6.05/7	
ZmMLO8 9.04	9.05 Southwestern Corn Borer QTL
ZmMLO9 na	

TABLE 4

Mutator Insertion Mutants of ZmMLO Homologues (TUSC)

5	ZmMLO2	
	PV03 81D3	C. heterostrophus intermediate resistance (segregating as recessive)
10	PV03 81D2 (sib)	C. heterostrophus intermediate resistance (segregating as recessive)
	PV03 108B1	C. heterostrophus intermediate resistance (segregating as recessive)
	PV03 108A1 (sib)	C. heterostrophus resistance (segregating as recessive)
15	BT94 5F11	C. heterostrophus resistance (segregating as recessive); also leaf necrosis

The Trait Utility System for corn (TUSC) system was utilized to generate transposon mutants in the *Mlo* sequences in maize. The combination of transposon mutagenesis via *Mutator* and PCR-based selection of target-specific *Mu* insertions makes up the TUSC system. See, for example, Benson *et al.* (1995) *Plant Cell*. 7:75-84; Mena *et al.* (1996) *Science* 274:1537-1540; U.S. Patent Application No. 08/835,638 and U.S. Patent Application No. 08/262,056; all of which are herein incorporated by reference. Table 4 shows *Mutator* insertion mutants of *ZmMlo* homologues.

Oligonucleotide primers based on *Mlo* maize sequences were designed and paired with *Mu* TIR primer in PCR reactions. The products of the screening process identify individuals in the collection having *Mu* elements inserted into *Mlo* sequences.

The gene-specific primers used for TUSC of *ZmMLO2* are: 19005, 5'-AGTTATCCAAGGCATGCCGGTGGTAAACT-3' (SEQ ID NO:18); and 19007, 5'-AAGTCATCGAGGGAGGCATCACTTACAG-3' (SEQ ID NO:19). For TUSC the mutator-element specific primer used was: 9242, 5'-AGAGAAGCCAACGCCA(AT)CGCCTC(CT)ATTCGTC-3' (SEQ ID NO:20). Seven positive "hits" to *ZmMLO2* were identified. Twenty F2 seeds for each of the hits were planted in the greenhouse and screened as follows. Seeds were planted in trays, containing 2-inch wells for each seed, in Strong-Lite (circleR) Universal Mix potting soil (Universal Mix, Pine Bluff, Arkansas, U.S.A.) and grown in a greenhouse (16 hour day, 20.6-34.8°C, 50% average RH, 560-620 microEinsteins, of light both sun and halogen lamps, averaging 25 heat units per day. Plants were grown for 14-18 days until the V5 stage (ca. 25cm height) and inoculated with spores of *Cochliobolus heterostrophus* (Drechs.) Drechs. Race 0, also known as *Bipolaris maydis* (Southern Leaf Blight) and *Cochliobolus carbonum* R. R Nelson, also known as *Bipolaris zeicola* (Helminthosporium Leaf Spot). For general leaf inoculation, spore suspensions of 4×10^4 per ml of 0.02% Tween-20 were sprayed as an aerosol on the leaves. Approximately 0.5 ml were applied per leaf late in the afternoon. The plants were then immediately covered with a plastic tent and kept at R.T. in order to enhance humidity and spore germination. The plastic tent was removed early in the morning, and the plants were returned to the greenhouse for the duration of the experiment, usually three days. Susceptibility to *Cochliobolus carbonum* was determined by either increased size or number of lesions. Resistance to *Cochliobolus*

heterostrophus was determined by either reduced size or number of lesions. Lesion mimics were also investigated; these mutations form spontaneous lesions in the absence of pathogen attach. Other phenotypes affecting growth and development were also noted.

TABLE 5

Maize <i>MLO</i> Gene	ZmMLO1	ZmMLO2	ZmMLO4	ZmMLO5	ZmMLO6a	ZmMLO6b	ZmMLO7	ZmMLO8	ZmMLO9
Gene Mapping Information									
Gene Complexity on Southern blot	1	1	1	1	2	2	1	1, 2	1, 2
Chromosome location (bin)	1.01	1.04	3.05	3.06	5.05	6.02/3	9.04	6.05/7	unassigned
Chromosome location (arm)	1S	1S	3L	3L	5L	6L	9L	6L	
Nearby Public Marker	BnI5.62	BnI12.06	umc10	bnl10.24	umc43	umc113	umc95	bnl5.47	
Distance from Public Marker	12.3 (down)	10.1 (down)	6.5 (down)	1.3 (up)	10.9 (down)	52.0 (up)	6.3 (down)	10.1 (up)	
Distance	15	5.9	35.2	1.3	1.8	52	10.7	10.1	
LOD Score	18.2	24.1	21.3	31.7	35.2	4.6	36.5	23.8	
RFLP Restriction Enzyme	EcoRI	HindIII	EcoRV	EcoRI	BamHI	BamHI	HindIII	EcoRV	
Mapping Population	ALEB9	ALEB9	DRAG2	ALEB9	DRAG2	DRAG2	DRAG2	DRAG2	DRAG2
Mapping Parents (Public Names)	R67/P38	R67/P38	17J/MWO	R67/P38	17J/MWO	17J/MWO	17J/MWO	17J/MWO	17J/MWO

TABLE 6

Maize <i>MLO</i> Gene	Hv <i>MLO1</i>	Zm <i>MLO</i> 1	Zm <i>MLO2</i>	Zm <i>MLO4</i>	Zm <i>MLO5</i>	Zm <i>MLO6</i>	Zm <i>MLO7</i>	Zm <i>MLO8</i>	Zm <i>MLO9</i>
Protein Structure Predictions									
Number of amino acids	533	224+	565	509	149+	515	499	492	469+
Calculated molecular weight	60,408		64,274	58,613		56,980	56,806	55,500	
Calculated pI	9.7		9.6	9.3		9.5	8.9	9.3	
Number of transmembrane domains	6 or 7		6 or 7	6 or 7		6 or 7	6 or 7	6 or 7	
Glycosylation sites? Asn-X-Ser/Thr	0	0	1	3	2	1	1	0	
Signal Peptide	No	No	No	No	No	No	No	No	No

Example 1: Transformation and Regeneration of Transgenic Plants

Immature maize embryos from greenhouse donor plants are bombarded with a plasmid containing the *ZmMLO1* antisense sequence operably linked to the ubiquitin promoter (Figure 3). This plasmid also contains the selectable marker gene PAT (Wohlleben *et al.* (1988) *Gene* 70:25-37) that confers resistance to the herbicide Bialophos. Transformation is performed as follows. All media recipes are in the Appendix.

Preparation of Target Tissue

The ears are surface sterilized in 30% Chlorox bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are excised and placed embryo axis side down (scutellum side up), 25 embryos per plate, on 560Y medium for 4 hours and then aligned within the 2.5-cm target zone in preparation for bombardment.

Preparation of DNA

A plasmid vector comprising the PAT selectable marker and the *ZmMlo1* antisense sequence operably linked to a the ubiquitin is made and precipitated onto 1.1 μm (average diameter) tungsten pellets using a CaCl_2 precipitation procedure as follows:

- 100 μl prepared tungsten particles in water
- 10 μl (1 μg) DNA in TrisEDTA buffer (1 μg total)
- 100 μl 2.5 M CaCl_2
- 10 μl 0.1 M spermidine

Each reagent is added sequentially to the tungsten particle suspension, while maintained on the multitube vortexer. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid removed, washed with 500 ml 100% ethanol, and centrifuged for 30 seconds. Again the liquid is removed, and 105 μl 100% ethanol is added to the final tungsten particle pellet. For particle gun bombardment, the

tungsten/DNA particles are briefly sonicated and 10 µl spotted onto the center of each macrocarrier and allowed to dry about 2 minutes before bombardment.

Particle Gun Treatment

- 5 The sample plates are bombarded at level #4 in particle gun #HE34-1 or #HE34-2. All samples receive a single shot at 650 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

Subsequent Treatment

- 10 Following bombardment, the embryos are kept on 560Y medium for 2 days, then transferred to 560R selection medium containing 3 mg/liter Bialaphos, and subcultured every 2 weeks. After approximately 10 weeks of selection, selection-resistant callus clones are transferred to 288J medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos are transferred to
- 15 medium for germination and transferred to the lighted culture room. Approximately 7-10 days later, developing plantlets are transferred to 272V hormone-free medium in tubes for 7-10 days until plantlets are well established. Plants are then transferred to inserts in flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to classic
- 20 600 pots (1.6 gallon) and grown to maturity. Plants are monitored and scored for enhanced disease resistance as a result of antisense disruption of *ZmMlo1* function.

APPENDIX**272 V**

Ingredient	Amount	Unit
D-I H ₂ O	950.000	ml
MS Salts (GIBCO 11117-074)	4.300	G
Myo-Inositol	0.100	G
MS Vitamins Stock Solution ##	5.000	ml
Sucrose	40.000	G
Bacto-Agar @	6.000	G

Directions:

- 5 @ = Add after bringing up to volume

Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

- 10 ## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month. unless contamination or precipitation occurs, then make fresh stock.
- 15 Total Volume (L) = 1.00

288 J

Ingredient	Amount	Unit
D-I H ₂ O	950.000	ml
MS Salts	4.300	g
Myo-Inositol	0.100	g
MS Vitamins Stock Solution ##	5.000	ml
Zeatin .5mg/ml	1.000	ml
Sucrose	60.000	g
Gelrite @	3.000	g
Indoleacetic Acid 0.5 mg/ml #	2.000	ml
0.1mM Absciscic Acid	1.000	ml
Bialaphos 1mg/ml #	3.000	ml

Directions:

@ = Add after bringing up to volume

- 5 Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

Add 3.5g/L of Gelrite for cell biology.

- 10 ## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occurs, then make fresh stock.
- 15 Total Volume (L) = 1.00

560 R

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511	1.000	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	30.000	g
2, 4-D 0.5mg/ml	4.000	ml
Gelrite @	3.000	g
Silver Nitrate 2mg/ml #	0.425	ml
Bialaphos 1mg/ml #	3.000	ml

Directions:

- 5 @ = Add after bringing up to volume
 # = Add after sterilizing and cooling to temp.
 Dissolve ingredients in D-I H₂O in sequence
 Adjust to pH 5.8 with KOH
 Bring up to volume with D-I H₂O
- 10 Sterilize and cool to room temp.
 Total Volume (L) = 1.00

560 Y

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511	1.000	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	120.000	g
2,4-D 0.5mg/ml	2.000	ml
L-Proline	2.880	g
Gelrite @	2.000	g
Silver Nitrate 2mg/ml #	4.250	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H₂O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H₂O

Sterilize and cool to room temp.

10 ** Autoclave less time because of increased sucrose**

Total Volume (L) = 1.00


All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and
5 individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 5, lines 11-20	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) Manassa, Virginia USA	
Date of deposit	Accession Numbers 98725-98729
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indicators are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") Date of Deposit	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

WHAT IS CLAIMED IS:

1. A method for creating or enhancing disease resistance in a plant, said method comprising modulating the expression of an *Mlo* nucleotide sequence wherein said nucleotide sequence is selected from the group consisting of:

- (a) a nucleotide sequence encoding a maize *Mlo* gene;
- (b) a nucleotide sequence encoding at least one of the maize MLO amino acid sequences selected from the group consisting of the sequences set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, and 16;
- (c) a nucleotide sequence comprising at least one of the nucleotide sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15;
- (d) a nucleotide sequence selected from the group consisting of sequences deposited in the American Type Culture Collection as Accession Nos. 98725, 98726, 98727, 98728, and 98729;
- (e) a nucleotide sequence that encodes an antisense sequence corresponding to a sequence of a), b), c) or d);
- (f) a nucleotide sequence which hybridizes under stringent conditions to one of the nucleotide sequences set forth in a), b), c), d) or e).

2. The method of claim 1, wherein said modulation comprises decreasing the expression of the *Mlo* sequence.

3. The method of claim 1, wherein said plant is a monocot.

4. The method of claim 3, wherein said monocot is maize.

5. A modified plant having altered levels of an MLO protein wherein said plant has stably incorporated into its genome a DNA construct, wherein said DNA construct comprises a nucleotide sequence operably linked

to a promoter capable of regulating transcription of said sequence in said plant, wherein said nucleotide sequence is selected from the group consisting of:

- (a) a nucleotide sequence encoding a maize *Mlo* gene;
- (b) a nucleotide sequence encoding at least one of the maize MLO amino acid sequences selected from the group consisting of the sequences set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, and 16;
- (c) a nucleotide sequence comprising at least one of the nucleotide sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15;
- (d) a nucleotide sequence selected from the group consisting of sequences deposited in the American Type Culture Collection as Accession Nos. 98725, 98726, 98727, 98728, and 98729;
- (e) a nucleotide sequence that encodes an antisense sequence corresponding to a sequence of a), b), c) or d);
- (f) a nucleotide sequence which hybridizes under stringent conditions to one of the nucleotide sequences set forth in a), b), c), d) or e).

6. The modified plant of claim 5, wherein said plant exhibits a decrease in the expression of a *Mlo* gene.

7. The modified plant of claim 5, wherein said plant is a monocot.

8. The modified plant of claim 7, wherein said monocot is maize.

9. The maize plant of claim 8, wherein said plant has decreased levels of the *Mlo* protein.

10. The maize plant of claim 8, wherein said maize plant comprises an insertion sequence in at least one maize *Mlo* gene.

11. Seed of the plant of any one of claims 5, 6, 7, 8, 9, and 10.

12. An isolated MLO protein, wherein said protein comprises an amino acid sequence selected from the group consisting of the MLO amino acid sequences set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, and 16.

13. An isolated nucleotide sequence said sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding a maize *Mlo* gene;
- (b) a nucleotide sequence encoding at least one of the maize MLO amino acid sequences selected from the group consisting of the sequences set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, and 16;
- (c) a nucleotide sequence comprising at least one of the nucleotide sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15;
- (d) a nucleotide sequence selected from the group consisting of sequences deposited in the American Type Culture Collection as Accession Nos. 98725, 98726, 98727, 98728, and 98729;
- (e) a nucleotide sequence that encodes an antisense sequence corresponding to a sequence of a), b), c) or d);
- (f) a nucleotide sequence which hybridizes under stringent conditions to one of the nucleotide sequences set forth in a), b), c), d) or e).

14. A modified plant cell having an altered level of an MLO protein wherein said plant cell has stably incorporated into its genome a DNA construct, wherein said DNA construct comprises a nucleotide sequence operably linked to a promoter capable of regulating transcription of said sequence in said plant cell, wherein said nucleotide sequence is selected from the group consisting of :

- (a) a nucleotide sequence encoding a maize *Mlo* gene;
- (b) a nucleotide sequence encoding at least one of the maize MLO amino acid sequences selected from the group consisting of the sequences set forth in SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, and 16;
- (c) a nucleotide sequence comprising at least one of the nucleotide sequences set forth in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, and 15;

- (d) a nucleotide sequence selected from the group consisting of sequences deposited in the American Type Culture Collection as Accession Nos. 98725, 98726, 98727, 98728, and 98729;
- (e) a nucleotide sequence that encodes an antisense sequence corresponding to a sequence of a), b), c) or d);
- (f) a nucleotide sequence which hybridizes under stringent conditions to one of the nucleotide sequences set forth in a), b), c), d) or e).

15. The modified plant cell of claim 14, wherein said cell exhibits a decrease in the expression of an *Mlo* gene.

16. The modified plant cell of claim 14, wherein said cell is from a monocot.

17. The modified plant cell of claim 16, wherein said monocot is maize.

1/5

CLUSTAL W (1.7) Multiple Sequence Alignments of HvMLO1 and 9 ZmMLO Homologues

```

HvMLO1 -MSDKKGVPAELPETPSWAVAVVFAAMVLVSVLMEHGLHKLGHWFQHRHKKALWEALEK
ZmMLO1 -----
ZmMLO2 MGGDG----TRALDQPTWAVAACVAVIVAASILLEGLHHLGQLLNKKRKKALFDALEK
ZmMLO6 MGGGGGGGNSRELDQPTWAVASVCGVIVLISILLEKGLHHVGEFFSHRKKKAMVEALEK

ZmMLO7 -----MGKEATLAFTPTWVVAIVCLVIVSISLAAERSLHYLGKYLECKKQKALFSALQR
ZmMLO5 -----
ZmMLO8 ---MAAEGEAAALEFTPTWIVAACSLIVLLSLVAERCLHYLGKTLKRKNQKPLFEALLK
ZmMLO9 -----
ZmMLO4 ----MAAEQGRSLAETPTWSVATVTTLMVAAACFLVERSLSRFAKWLRTKRKAMLALEK
Regions[---N-Terminus---] [---TM Helix 1---] [---Intervening Region 1---]

HvMLO1 MKAELMLVGFISLLLIVTQDPIIAKICISED-AADVMWPCKRGTEGRKPS-----
ZmMLO1 -----
ZmMLO2 VKSEMLTLGFISLLLIVTGR-YIARICIEPG-AANTMLPCRLSGHSAECPKGHGRR---
ZmMLO6 VKAELMLVGFISLLLVFGQN-YIIVKVCISNH-AANTMLPCKLEAAAVEGK-DGHGKEAAA

ZmMLO7 LKEELMLLGFISFVLSLSQG-FIVSICIPET-STDFMLPCNR-----GNS-----
ZmMLO5 -----
ZmMLO8 VKEELMLLGFISLLLVTFQG-MIRRTCIPER-WTFHMLPCEKPEKAGEAATMEHFVGLT
ZmMLO9 -----
ZmMLO4 IREELMLLGVISLLLSQTAR-FISEICVPSSLFTSRFYICESDYQDLLRNTDANQTALD
---] [-----TM Helix 2-----] [-----Intervening Region 2-----]

HvMLO1 -----K-----YVDYCEPGKVA
ZmMLO1 -----
ZmMLO2 -----HLS-----EDPTNLFSCRKGMVS
ZmMLO6 VVAGKKKVAVAVPGKKKKKAAAADHLGGVVDWPPYYAHNARMLAEASMATKCEPGKVP

ZmMLO7 -----RVAEGAKICNKKGDVP
ZmMLO5 -----
ZmMLO8 GRIGRR-----LLQEGTAGAEQCQKKGKVP
ZmMLO9 -----HSPRRNLKTSYHNHEGCREGYES
ZmMLO4 -----KNMFGGQ-RLHVCGEHHEP
-----Intervening Region 2-----

HvMLO1 LMSTGSLHQLHVFIFVLAVFHVTYSVITIALSRLKMRTWKKWETETTSLEYQFANDPAR-
ZmMLO1 -----
ZmMLO2 LVSADGMHQLHIFVFFLAVFHVTFSTFMSLGRAKTRIKVWEKETCSPQYNYLNDPSK-
ZmMLO6 LISINALHQLHIFIFVLAVFHVSYSAITMALGRAKIRAWKEWEKAAGQDYEFSDPTR-

ZmMLO7 LLSLEALHQLHIFIFVLGLVHVVECATTSISFGAKMRKWKHWETEIHREVHEKLQGEKNE
ZmMLO5 -----
ZmMLO8 LLSLEAIHQLHIFIFVLAITHVIFSVTTMLLGGQIHQWKQWENGIKKDAPGNGPKVTN-
ZmMLO9 FVSHEGLEQLHRFIFVMVTHVTYSCLTMLLAILKHKWRDWEDEAFRDNHESFSQIAYE
ZmMLO4 FVSYEGLEQLHRFLFILGITHVLYSFVTVVLMSIKITYSWRKWETLAGPIAAEEEEKARRT-
-----] P---TM Helix 3---] [-----Intervening Region 3-----]

HvMLO1 -----FRFTHQTSFVKRHL--GLSSTPGIRWVVAFFRQFFRSVTKVDYLTLAGF
ZmMLO1 -----
ZmMLO2 -----FRLTHQTSFVRGHAS-CWSKSTITLYFVSFFRQFFRSVRKTDYFTLRHGF
ZmMLO6 -----FRFTHQTSFVRQHNM-VLNKFPASFYISNFFRQFFRSVRQADYCALRHSF

ZmMLO7 GQGSSLSIVVLHREHQDEFVHKRTKGFWMKLAVASWITAFKQFQDSVSKSDYEALRSAF
ZmMLO5 -----
ZmMLO8 -----VHHHEFIKKRFGIGKDSIILSWLHSFGKQFYRSVSKSDYTMRLGF
ZmMLO9 S-----ATRRQPALTKSYSFRSWSQNNVVMWLVCFIAQFGQSVVRADYLILRKGF
ZmMLO4 -----KVMRRQSTFVFNASHPWSKNKILIWMLCFLRQFKGSIIRSDYLALRLGF
-----Intervening Region 3-----

```

TO FIG. 1B.

FIG. 1A.

2/5
FROM FIG. 1A.

HvMLO1 INAHLSONSKFDFHKYIKRSMEDDFKVVVGISLPLWGVAILTLFLDINGVGTLIWISFIP
ZmMLO1 -----
ZmMLO2 ISAHLSPGTRFNFRKYIKRSLEDDFKTVVGISPLWASALAVMLFNVHGWHLFWFSAIP
ZmMLO6 VNVHLAPGSKFDFDKYIDRSLEDDFKVIVGISPLWASALIFLFLNVNGWHTMLWISIMP
ZmMLO7 VVEHYEKPDI DFHKYMTAVEYEFKRVVGISWYLVLFVILFLLNINNGWHTYFWLAFIP
ZmMLO5 -----
ZmMLO8 IMTHCPGNPKFDFHRYMVRVLEADFKKVVGISWYLVFVIFLLNINNGWHTYFWIAFLP
ZmMLO9 IMTHNLPPT-YDFHNYMIRSMEEFEKIVGVSGLLWGFVFAFMFNVDGSNLYFWIAILP
ZmMLO4 MTYHKLPHS-YDFHKYMVRSMEDDYNIGTIGISWPLWAYAIVCILINVHGINIYFWLSFVP
---Intervening Region 3---] [----TM Helix 4---] [IR 4] [-----TM
HvMLO1 LVILLCVGKLEMIIMEMALEIQDRAS--VIKGAPVVEPSNKFVWHRPDVVLFFIHLTL
ZmMLO1 -----EIQDKAT--VIKGAPVVEPSDRFFWFNRPQWVLFILHLTL
ZmMLO2 LVVILAVGKLAIAIMMAIEIAERHT--VIQGMPPVKLSDDHFWFGKPRVLVHLIHFAS
ZmMLO6 VVILSVGKLGICRMAIDITERHA--VIQGI PMVQVSDSYFWFARPTFVLFIHFTL
ZmMLO7 LFLLLIVGAKLEGIITRLAQEAASLSN-NTEEVPKIKPCKDHFVWFKPELVHLIHFIL
ZmMLO5 -----IHFIL
ZmMLO8 LILLIAIGTKLEHVIAQLAHDVAEKHT--AVEGDVIVKPSDEHFWFGKPRVILYLIHFIL
ZmMLO9 VTLVLLVGAKLQHVIAITLTAEGAKMST-----YGPRIQPRDDLFWFKKPEFLLWL IHFVL
ZmMLO4 VELVLLVGTELQHVIAQLALEVAEATA--PY-VGSQLKLRDDLFWFGKPRVLWVLIQFIS
Helix 5] [-----Intervening Region 5-----
HvMLO1 FQNAFQMAHFVWTVATPGLKKCYHTQIGLSIMKVVVGLALQFLCSYMTFPLYALVTQMGs
ZmMLO1 FQNAFQMAHFVWTLTPDLKKCYHERLGLSIMKVAVGLVLQVLCYITFPLYALVTQMGs
ZmMLO2 FQNAFEITYFFWIWYEFGLRSCFHDNFEFIIARVCLGATVQFMCSYITLPLAYALVSQMGs
ZmMLO6 FQNGFQIIYFLWILYEGMDSCFNDSEEFVFA RLCLGVVVQVLC SYVTLPYALVSQMGs
ZmMLO7 FQNSFEISFFFWILVSEGGFGSCMMERKPYVISRLVIGVIIIEVECSYITLPLYAIVTHMTG
ZmMLO5 FQNAFEIAFFFWILT TYGFNSCIMDHVPFIVPRLVVGAIQLLCSYSTLPYAI VTQMGs
ZmMLO8 FQNAFEIAFFFWILSTYGFDS CIMGQVRFIVPRLVIGVVIQLLCSYSTLPYAI VTQMGs
ZmMLO9 GQNAFELASFFWFWQFGYDSCFIKNHLLVYCRLILGFAGQFLCSYSTLPVYALVTQMGs
ZmMLO4 FQNAFELATFLWSLWELSAQTCFMKHYYMVAIRLISGLLVQFWCLYYSTLP LNVIISQMGs
-----Intervening Region 4-----] [-----TM Helix6-----] [---
HvMLO1 NMKRSIEDQTSKALTNWRNTAKEKKKVR--DTDM LMAQMIGDATPSRGSS--PMPSRGSS
ZmMLO1 HMKKTIFEEQTAKAVMKWRKTAKDKVRQRE--AAGFLDVLT SADTTPSHSRA--TSPSRGNS
ZmMLO2 EMKRTIFDEQTAKALKKWKHKA VVKKKHKK-----DSSHNSSETPSTDTTG PAGEAGEW
ZmMLO6 TMKQSFDEQTSKALKNWRAGAKKKAPTG-----GSKHGGGGSPTAGGS--PTKADGDA
ZmMLO7 QIKLHGFGSRVHESVHGWIGLRKKPF SFWKIPGGDPNADSGREADVTRRVAKERSGSSRS
ZmMLO5 FFKKEIFDEHVQQSLLGWAQKAKKRKALRNNGNGSNGAAAG--SSHPSATARLELMRRAVA
ZmMLO8 CYKKEIFNEHVQQGLGWAQKV KMKKGLK--G-----AAS--ASKDESITNADSAGPSVK
ZmMLO9 KYKAALIPRIRET IHGWGKATRKRRRR--RGDDSTVRTETSTVCSLTDDEDFEDDDD
ZmMLO4 KFKKSLVSENVRESLSWCKRVKDRSRHN--PLFSRNGLTTRSVCSLDTTYETDHTNT
-----C-Terminus-----
HvMLO1 -----PVHLLHKGM-----GRSDDPQSAPTSPT-----QQEARDMYPVVVAH
ZmMLO1 -----PVHLLHKYR-----GRSEEPQSGPASP-----GRELDMYPVADQH
ZmMLO2 QRLHEVPVRHLHRYKTI-----AHVGGVRSPLSDPDYSDDTEPLSLQTRHLIPPAKQR
ZmMLO6 -----
ZmMLO7 -----MPMAPADEI-----VTVDVAVAAAAVGGGP-----
ZmMLO5 -----LEEGSAGGN-----GSEASAAELHDTGPKL-----
ZmMLO8 -----IEMAKAG-----EDVEIVGNTG-----
ZmMLO9 HHHHGPSYDTPRAGGRPPYKLIETHRQSGSGHDGPRPGGTPCFHPSGSGSGHAMLLRQAS
ZmMLO4 VCTLSRTASATSLDDQL-----TVVTVDDEPSCIEKDV-----
-----C-Terminus-----

TO FIG. 1C.

FIG. 1B.


3/5

FROM FIG. 1B.

HvML01	PVHRLNPNDRRRSASSSALEADIPSADF SFSQG-----	533 aa
ZmML01	RLHRLDP-ERM RPASSTAVNIDIADDF SFSMR-----	224 aa
ZmML02	SLDTERAEVRVNVVETAAAPSDVLQDSFSFPRLLP RHVPDK--	565 aa
ZmML06	-----	515 aa
ZmML07	-----	499 aa
ZmML05	-----	149 aa
ZmML08	-----	492 aa
ZmML09	VSAPSSPSYRGGNNVTRSASMPGIAALRTTGSGTPTRVSHEEPT	469 aa
ZmML04	-----	509 aa
	-----C-Terminus-----	

FIG. 1C.

FIG. 2.



5/5




FIG. 3.

SEQUENCE LISTING

<110> Briggs, Steven

<120> Manipulation of Mlo Genes to Enhance Disease Resistance
in Plants

<130> 5718-42035718/158714

<140>

<141>

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 845

<212> DNA

<213> Zea mays

<220>

<223> Mlo1

<400> 1

```
aggagataca ggacaaggcg acgggtcatca aggggggcgcc tgtggtggag ccaagtgaca 60
ggttctttctg gtttaaccgc cctggctggg tcctcttctt catccacctc acgtctttcc 120
agaacgcctt ccagatggcg catttcgttt ggacactgct caccacagac ctgaagaaat 180
gctaccacga gaggtctggc ctgagcatca tgaaagttgc ggtggggctg gttctccagg 240
tcctctgcag ctacatcacc ttcccgtctt acgcgtctgt cacgcagatg gggtcgcaca 300
tgaagaagac catcttcgag gagcagacgg ccaaggcggt gatgaagtgg cgcaagacgg 360
ccaaggataa ggtgcggcag cgggaggcgg caggcttctt cgacgtgctg acgagcgccg 420
acaccacgcc gagccacagc cgcgcgacgt cgccgagccg gggcaactcg ccggtgcacc 480
tgctccacaa gtacaggggc aggtcggagg aaccgcagag cgggccggcg tcgccggggc 540
gggagctcgg ggacatgtac ccggtggctg accagcatcg cctgcacagg ctggaccccc 600
agaggatgag gccgcctcg tccaccgccg tcaacattga catcgctgat gccgattttt 660
cttttagcat gcggtgacct gaccttgaac gaattctgtg tccttactct tgtataggga 720
agcaaaagca tagacggaga acataatgac acgttacgtt agggaaaagt tcgtttattc 780
atcataaaat agaatacgta ataaactagt atctccctct tcaaaaaaaaa aaaaaaaaaa 840
aaaaa
```

845

<210> 2

<211> 224

<212> PRT

<213> Zea mays

<220>

<223> Mlo1

<400> 2

Glu Ile Gln Asp Lys Ala Thr Val Ile Lys Gly Ala Pro Val Val Glu
 1 5 10 15

Pro Ser Asp Arg Phe Phe Trp Phe Asn Arg Pro Gly Trp Val Leu Phe
 20 25 30

Leu Ile His Leu Thr Leu Phe Gln Asn Ala Phe Gln Met Ala His Phe
 35 40 45

Val Trp Thr Leu Leu Thr Pro Asp Leu Lys Lys Cys Tyr His Glu Arg
 50 55 60

Leu Gly Leu Ser Ile Met Lys Val Ala Val Gly Leu Val Leu Gln Val
 65 70 75 80

Leu Cys Ser Tyr Ile Thr Phe Pro Leu Tyr Ala Leu Val Thr Gln Met
 85 90 95

Gly Ser His Met Lys Lys Thr Ile Phe Glu Glu Gln Thr Ala Lys Ala
 100 105 110

Val Met Lys Trp Arg Lys Thr Ala Lys Asp Lys Val Arg Gln Arg Glu
 115 120 125

Ala Ala Gly Phe Leu Asp Val Leu Thr Ser Ala Asp Thr Thr Pro Ser
 130 135 140

His Ser Arg Ala Thr Ser Pro Ser Arg Gly Asn Ser Pro Val His Leu
 145 150 155 160

Leu His Lys Tyr Arg Gly Arg Ser Glu Glu Pro Gln Ser Gly Pro Ala
 165 170 175

Ser Pro Gly Arg Glu Leu Gly Asp Met Tyr Pro Val Ala Asp Gln His
 180 185 190

Arg Leu His Arg Leu Asp Pro Glu Arg Met Arg Pro Ala Ser Ser Thr
 195 200 205

Ala Val Asn Ile Asp Ile Ala Asp Ala Asp Phe Ser Phe Ser Met Arg
 210 215 220

<210> 3
 <211> 2030
 <212> DNA
 <213> Zea mays

<220>
 <223> M1o2

<400> 3
 gcggcaagca attgttcat ctgtccact ccattccat ttaacgcact ctccactcct 60
 tcgcggcggg ggtttaggcg cacggcggca ggccaattaa cgcgctgcg gcgagcaccg 120
 ggggggaaat gggcggcgac ggcacgagg cgctggacca gacgccgacg tgggccgtgg 180
 ccgccgtctg cgtgtctatc gtcgcggcgt ccatactcct cgagggcttc ctccaccacc 240
 tcggtcagtt gctcaacaag aagcggaga aggcgctgtt tgacgctctg gagaagggtta 300
 aatcgagct gatgactctg gggttcatat cgctgctgct gaccgtgacg ggaagggtaca 360
 tcgcgcgcat ctgcatcccg gagggagccg caaacaccat gctgccctgc cgctgtccg 420
 gtcactcggg ggcggaagag cccaagggcc atggccgacg gcacctatct gaagatccaa 480
 ccaacctttt ttctgcccgc aaaggcatgg tgcgctctgt ttcagccgac ggcattgcat 540
 agctgcacat ttctgtgttc ttcttgccg tcttccatgt taccttcagt ttcttcacaa 600
 tgtccttggg tagagcaaag actcgtatat ggaagggtgt ggaaaaggaa acttgctccc 660
 ctgagtataa ttatttaaat gacctctcaa agttcaggct tacgcaccaa acatcttttg 720
 tgaggcaaca tgcaagttgt tggagcaaaa gcacaatcac gctctatatt gtgagcttct 780
 ttaggcagtt ctccagatcc gtccgtaaga cagactactt tactttgcga catggattca 840
 tttcagctca tttatctccg gggactaggt tcaattttcg aaagtacatt aaaaggctct 900
 tggaggatga tttcaagaca gttgttgga ttagtccacc actatgggct tctgcttttg 960
 ctgtcatgct gttcaatggt catggatggc acaacttatt ctggttctct gcaattcccc 1020
 ttgtagttat tctagcagtt ggaacaaagc tgcaggctat aattgctatg atggctattg 1080
 aaattgcaga gaggcataca gttatccaag gcatgccggg ggtaaaacta agtgatgatc 1140
 atttctggtt cgggaagcct cgtttggttc tccacctcat tcatttcgca tcatttcaga 1200
 acgcatttga aattacatat ttcttttgga tctggtacga atttggcctg aggtcctgct 1260
 tccatgacaa ctttgagttt atcattgcaa gagtctgcct tggggctatc gtccagttta 1320
 tgtgcagcta catcaccctt cactctatg cccttgtttc tcagatgggt tcagaaatga 1380
 agcgaacaat ttctgacgag cagacagcga aggcctgaa gaaatggcac aaggccgtgg 1440
 tgaagaagaa acaccacaag gattcctcac acaattcttc cgagactcct agcactgaca 1500
 caacaggacc tgcaggggaa gcgggtgagt ggcagcggct gcacgaggtg ccggctccggc 1560
 acctccaccg gtacaagacc atcgcgcacg tcggcggcgt gaggagcccg ttgtcggacc 1620
 cggactacag cgacacggac gacaccgagc cgctgtcgtt gcagactagg cacctgatac 1680
 cgccggcgaa gcagcggagc ctgcacaccg agcgcgcgga ggtgcgcgtg aacgtcgtag 1740
 agacagcggc ggcgccaaag gacgtcctcc aagacagctt ctcgttccca aggtcgttc 1800
 ctctcgcga tgtgccggac aagtaaagtg gccagcagtc tgcttagtac tacagactgt 1860
 cctatataga gatagtagaa tagtagtttc aagttataag tcgatttggc tagttataca 1920
 acttttgcta ttattttaga taaaggctat gtctatatag caaaagtgtg aaaactagaa 1980
 aaatcaaaat gatttataat ttgaaaaaaa aaaaaaaaaa aaaaaaaaaa 2030

<210> 4
 <211> 565
 <212> PRT
 <213> Zea mays

<220>

<223> M1o2

<400> 4

```

Met Gly Gly Asp Gly Thr Arg Ala Leu Asp Gln Thr Pro Thr Trp Ala
  1           5           10           15

Val Ala Ala Val Cys Ala Val Ile Val Ala Ala Ser Ile Leu Leu Glu
          20           25           30

Gly Phe Leu His His Leu Gly Gln Leu Leu Asn Lys Lys Arg Lys Lys
          35           40           45

Ala Leu Phe Asp Ala Leu Glu Lys Val Lys Ser Glu Leu Met Thr Leu
          50           55           60

Gly Phe Ile Ser Leu Leu Leu Thr Val Thr Gly Arg Tyr Ile Ala Arg
          65           70           75           80

Ile Cys Ile Pro Glu Gly Ala Ala Asn Thr Met Leu Pro Cys Arg Leu
          85           90           95

Ser Gly His Ser Val Ala Glu Glu Pro Lys Gly His Gly Arg Arg His
          100          105          110

Leu Ser Glu Asp Pro Thr Asn Leu Phe Ser Cys Arg Lys Gly Met Val
          115          120          125

Ser Leu Val Ser Ala Asp Gly Met His Gln Leu His Ile Phe Val Phe
          130          135          140

Phe Leu Ala Val Phe His Val Thr Phe Ser Phe Phe Thr Met Ser Leu
          145          150          155          160

Gly Arg Ala Lys Thr Arg Ile Trp Lys Val Trp Glu Lys Glu Thr Cys
          165          170          175

Ser Pro Gln Tyr Asn Tyr Leu Asn Asp Pro Ser Lys Phe Arg Leu Thr
          180          185          190

His Gln Thr Ser Phe Val Arg Gln His Ala Ser Cys Trp Ser Lys Ser
          195          200          205

Thr Ile Thr Leu Tyr Phe Val Ser Phe Phe Arg Gln Phe Phe Arg Ser
          210          215          220

Val Arg Lys Thr Asp Tyr Phe Thr Leu Arg His Gly Phe Ile Ser Ala

```


225					230						235				240
His	Leu	Ser	Pro	Gly	Thr	Arg	Phe	Asn	Phe	Arg	Lys	Tyr	Ile	Lys	Arg
				245					250					255	
Ser	Leu	Glu	Asp	Asp	Phe	Lys	Thr	Val	Val	Gly	Ile	Ser	Pro	Pro	Leu
			260					265					270		
Trp	Ala	Ser	Ala	Leu	Ala	Val	Met	Leu	Phe	Asn	Val	His	Gly	Trp	His
	275						280					285			
Asn	Leu	Phe	Trp	Phe	Ser	Ala	Ile	Pro	Leu	Val	Val	Ile	Leu	Ala	Val
	290					295					300				
Gly	Thr	Lys	Leu	Gln	Ala	Ile	Ile	Ala	Met	Met	Ala	Ile	Glu	Ile	Ala
305					310					315					320
Glu	Arg	His	Thr	Val	Ile	Gln	Gly	Met	Pro	Val	Val	Lys	Leu	Ser	Asp
			325					330					335		
Asp	His	Phe	Trp	Phe	Gly	Lys	Pro	Arg	Leu	Val	Leu	His	Leu	Ile	His
			340					345					350		
Phe	Ala	Ser	Phe	Gln	Asn	Ala	Phe	Glu	Ile	Thr	Tyr	Phe	Phe	Trp	Ile
		355					360					365			
Trp	Tyr	Glu	Phe	Gly	Leu	Arg	Ser	Cys	Phe	His	Asp	Asn	Phe	Glu	Phe
	370					375					380				
Ile	Ile	Ala	Arg	Val	Cys	Leu	Gly	Ala	Ile	Val	Gln	Phe	Met	Cys	Ser
385					390					395					400
Tyr	Ile	Thr	Leu	Pro	Leu	Tyr	Ala	Leu	Val	Ser	Gln	Met	Gly	Ser	Glu
			405						410				415		
Met	Lys	Arg	Thr	Ile	Phe	Asp	Glu	Gln	Thr	Ala	Lys	Ala	Leu	Lys	Lys
			420					425					430		
Trp	His	Lys	Ala	Val	Val	Lys	Lys	Lys	His	His	Lys	Asp	Ser	Ser	His
		435					440					445			
Asn	Ser	Ser	Glu	Thr	Pro	Ser	Thr	Asp	Thr	Thr	Gly	Pro	Ala	Gly	Glu
	450					455					460				
Ala	Gly	Glu	Trp	Gln	Arg	Leu	His	Glu	Val	Pro	Val	Arg	His	Leu	His
465					470					475					480
Arg	Tyr	Lys	Thr	Ile	Ala	His	Val	Gly	Gly	Val	Arg	Ser	Pro	Leu	Ser

485 490 495
 Asp Pro Asp Tyr Ser Asp Thr Asp Asp Thr Glu Pro Leu Ser Leu Gln
 500 505 510
 Thr Arg His Leu Ile Pro Pro Ala Lys Gln Arg Ser Leu Asp Thr Glu
 515 520 525
 Arg Ala Glu Val Arg Val Asn Val Val Glu Thr Ala Ala Ala Pro Ser
 530 535 540
 Asp Val Leu Gln Asp Ser Phe Ser Phe Pro Arg Leu Leu Pro Pro Arg
 545 550 555 560
 His Val Pro Asp Lys
 565

 <210> 5
 <211> 1841
 <212> DNA
 <213> Zea mays

 <220>
 <223> M1o4

 <400> 5
 agatatcagg gcagctggaa ctgaagggtgg cgggagccga gtgatccggc ggtgagctga 60
 gcggggcggg atggcgggcg agcaggggag gtcgctggcg gagacgcca cctgggtccgt 120
 ggcaaccgtc accacgctca tggctcgctgc ctgcttcctc gtcgagcgct ccctctcgcg 180
 cttcgccaag tggctgcgca agaccaagcg gaaggccatg ctgcccgcgc tcgagaagat 240
 ccgcaagag ctgatgctgc tcggagtcac ctcgctgctg ctcagccaga cggcgcgctt 300
 catatcggag atctgcgtgc cgtcctcgct cttcaccagc cgcttctaca tctgctccga 360
 gagcgactac caggacctgc tgcgcaaac ggacgccaac cagacggcgc tcgacaagaa 420
 catgttcggt ggccaacggc tgcacgtctg tggcgagggc catgaacctt ttgtttcgta 480
 cgagggccctt gagcagctgc accggtttct cttcatcctt ggtatcactc atgtgttgta 540
 cagttttgta acagtgggtc tgtccatgat caagatctat agctggagga agtgggaaac 600
 cttagcaggt ccaattgctg ctgaggaatt gaaagctagg agaaccaagg tgatgagaag 660
 gcagtcaacc tttgttttta acaatgcttc tcatccatgg agcaaaaata aaatacttat 720
 ttggatgctt tgctttttgc gtcaattcaa gggctccata ataaggtcag actatttggc 780
 actgagggtt ggctttgtca catatcacia gctaccacat tcatatgact tccataaata 840
 catggtacgg agcatggaag atgattacaa tgggactatt ggtatcagtt ggccactttg 900
 ggcatatgag attgtctgca tattaatcaa tgttcatggt atcaatatat atttctgggt 960
 gtcctttgtt cctgttattc tgggtgcttct agtgggtact gaacttcagc acgtcattgc 1020
 tcagttggct ttggaagtcg ctgaggcaac agcgccttat gttgggtcac aacttaaact 1080
 gcgtgatgat ctattttggt ttggaaagcc tcgggtactc tgggtggctta tacagttcat 1140
 ttcatttcag aatgcttttg agctggcaac attcttatgg tctctgtggg aactcagtcg 1200
 acaaacatgt ttcatgaagc actactacat gggttgccatt cggttgatgt ctgggctcct 1260

```

agttcagttt tgggtgcagct acagcacact gccgctgaat gtgattattt ctcagatggg 1320
ttccaagttc aagaaatcac tgggtctcgga gaacgtgagg gagtcgctgc acagctgggtg 1380
caagagggtt aaggacagga gccgacacaa tccgctcttc tcgcggaacg ggaccctcac 1440
gaccagatcc gtgtgctccc tagacaccac ctacgagacg gatcacgaga cgaacacggt 1500
gtgcacgctg tcgaggacgg cgtcggcgac gtcgctggac gaccagttga ccgtgggtcac 1560
cgtcgatgac gagccgtcct gcattgagaa ggatgtctga cgcagttgct gattcgccaa 1620
aactacatac tgcaccgacc tgtgtgttag gaggtactct aggactgaaa tgaatgtata 1680
tgtgcagtgg ttgcccctat cgtggcagtg accatgtgta tatatagaac taaaaaaaact 1740
ataatgtctc ttagctgaga gtgagaggct gtacatggaa atacttgagg cagcttgga 1800
caaatacaaa actgcattct ttcgagaaaa aaaaaaaaaa a 1841

```

<210> 6

<211> 509

<212> PRT

<213> Zea mays

<220>

<223> M104

<400> 6

```

Met Ala Ala Glu Gln Gly Arg Ser Leu Ala Glu Thr Pro Thr Trp Ser
  1              5              10              15

Val Ala Thr Val Thr Thr Leu Met Val Ala Ala Cys Phe Leu Val Glu
          20              25              30

Arg Ser Leu Ser Arg Phe Ala Lys Trp Leu Arg Lys Thr Lys Arg Lys
    35              40              45

Ala Met Leu Ala Ala Leu Glu Lys Ile Arg Glu Glu Leu Met Leu Leu
    50              55              60

Gly Val Ile Ser Leu Leu Leu Ser Gln Thr Ala Arg Phe Ile Ser Glu
    65              70              75              80

Ile Cys Val Pro Ser Ser Leu Phe Thr Ser Arg Phe Tyr Ile Cys Ser
          85              90              95

Glu Ser Asp Tyr Gln Asp Leu Leu Arg Asn Thr Asp Ala Asn Gln Thr
    100              105              110

Ala Leu Asp Lys Asn Met Phe Gly Gly Gln Arg Leu His Val Cys Gly
    115              120              125

Glu Gly His Glu Pro Phe Val Ser Tyr Glu Gly Leu Glu Gln Leu His
    130              135              140

Arg Phe Leu Phe Ile Leu Gly Ile Thr His Val Leu Tyr Ser Phe Val

```

145		150		155		160									
Thr	Val	Val	Leu	Ser	Met	Ile	Lys	Ile	Tyr	Ser	Trp	Arg	Lys	Trp	Glu
			165						170					175	
Thr	Leu	Ala	Gly	Pro	Ile	Ala	Ala	Glu	Glu	Leu	Lys	Ala	Arg	Arg	Thr
			180					185					190		
Lys	Val	Met	Arg	Arg	Gln	Ser	Thr	Phe	Val	Phe	Asn	Asn	Ala	Ser	His
		195					200				205				
Pro	Trp	Ser	Lys	Asn	Lys	Ile	Leu	Ile	Trp	Met	Leu	Cys	Phe	Leu	Arg
	210					215					220				
Gln	Phe	Lys	Gly	Ser	Ile	Ile	Arg	Ser	Asp	Tyr	Leu	Ala	Leu	Arg	Leu
225					230					235					240
Gly	Phe	Val	Thr	Tyr	His	Lys	Leu	Pro	His	Ser	Tyr	Asp	Phe	His	Lys
				245					250					255	
Tyr	Met	Val	Arg	Ser	Met	Glu	Asp	Asp	Tyr	Asn	Gly	Thr	Ile	Gly	Ile
			260					265					270		
Ser	Trp	Pro	Leu	Trp	Ala	Tyr	Ala	Ile	Val	Cys	Ile	Leu	Ile	Asn	Val
		275					280					285			
His	Gly	Ile	Asn	Ile	Tyr	Phe	Trp	Leu	Ser	Phe	Val	Pro	Val	Ile	Leu
	290					295					300				
Val	Leu	Leu	Val	Gly	Thr	Glu	Leu	Gln	His	Val	Ile	Ala	Gln	Leu	Ala
305					310					315					320
Leu	Glu	Val	Ala	Glu	Ala	Thr	Ala	Pro	Tyr	Val	Gly	Ser	Gln	Leu	Lys
				325					330					335	
Leu	Arg	Asp	Asp	Leu	Phe	Trp	Phe	Gly	Lys	Pro	Arg	Val	Leu	Trp	Trp
			340					345					350		
Leu	Ile	Gln	Phe	Ile	Ser	Phe	Gln	Asn	Ala	Phe	Glu	Leu	Ala	Thr	Phe
		355					360					365			
Leu	Trp	Ser	Leu	Trp	Glu	Leu	Ser	Ala	Gln	Thr	Cys	Phe	Met	Lys	His
	370					375					380				
Tyr	Tyr	Met	Val	Ala	Ile	Arg	Leu	Ile	Ser	Gly	Leu	Leu	Val	Gln	Phe
385					390					395					400
Trp	Cys	Leu	Tyr	Ser	Thr	Leu	Pro	Leu	Asn	Val	Ile	Ile	Ser	Gln	Met

405 410 415
 Gly Ser Lys Phe Lys Lys Ser Leu Val Ser Glu Asn Val Arg Glu Ser
 420 425 430
 Leu His Ser Trp Cys Lys Arg Val Lys Asp Arg Ser Arg His Asn Pro
 435 440 445
 Leu Phe Ser Arg Asn Gly Thr Leu Thr Thr Arg Ser Val Cys Ser Leu
 450 455 460
 Asp Thr Thr Tyr Glu Thr Asp His Glu Thr Asn Thr Val Cys Thr Leu
 465 470 475 480
 Ser Arg Thr Ala Ser Ala Thr Ser Leu Asp Asp Gln Leu Thr Val Val
 485 490 495
 Thr Val Asp Asp Glu Pro Ser Cys Ile Glu Lys Asp Val
 500 505

<210> 7

<211> 673

<212> DNA

<213> Zea mays

<220>

<223> M1o5

<400> 7

tgatccactt catcctcttc cagaacgcgt ttgagatcgc attcttcttc tggattctga 60
 ctacctacgg tttcaactcc tgcacatggt accacgtccc cttcattgtg ccgaggctcg 120
 ttgtcggggc catcatccag ctctctgtga gctacagcac cctgcctctg tacgcgattg 180
 tcacgcagat ggggactttc ttcaagaagg agatcttcga tgagcacgtg cagcagagcc 240
 ttctgggctg ggcgcagaag gccaaaga ggaaagcgct caggaacaat ggcaatggca 300
 gcaatggcgc cgcggccggg tcgtcacatc catctgcgac tgcgaggctc gaactgatga 360
 gacgagctgt cgctcttgaa gaaggcagtg ccggtggcaa cgggagtgag gctagtgcgg 420
 ccgagctaca cgacacgggc ccgaagctct gaacgctgta gaaaacgccc ggcagagcgt 480
 cagaagatgc agtgtttctg tacgcgagcc atatgtcact atgtcgtgtg gctgttttagc 540
 cagggtggtg agtgtgatac tgcgtatcat gatttgatg cttttctgga ttgagcttga 600
 agcgtgttat ggagtgggga gctttgtgta gtgttggtgt caaaaaaaaaa aaaaaaaaaa 660
 aaaaaaaaaa aaa 673

<210> 8

<211> 149

<212> PRT

<213> Zea mays

<220>

<223> Mlo5

<400> 8

Ile His Phe Ile Leu Phe Gln Asn Ala Phe Glu Ile Ala Phe Phe Phe
 1 5 10 15

Trp Ile Leu Thr Thr Tyr Gly Phe Asn Ser Cys Ile Met Asp His Val
 20 25 30

Pro Phe Ile Val Pro Arg Leu Val Val Gly Ala Ile Ile Gln Leu Leu
 35 40 45

Cys Ser Tyr Ser Thr Leu Pro Leu Tyr Ala Ile Val Thr Gln Met Gly
 50 55 60

Thr Phe Phe Lys Lys Glu Ile Phe Asp Glu His Val Gln Gln Ser Leu
 65 70 75 80

Leu Gly Trp Ala Gln Lys Ala Lys Lys Arg Lys Ala Leu Arg Asn Asn
 85 90 95

Gly Asn Gly Ser Asn Gly Ala Ala Ala Gly Ser Ser His Pro Ser Ala
 100 105 110

Thr Ala Arg Leu Glu Leu Met Arg Arg Ala Val Ala Leu Glu Glu Gly
 115 120 125

Ser Ala Gly Gly Asn Gly Ser Glu Ala Ser Ala Ala Glu Leu His Asp
 130 135 140

Thr Gly Pro Lys Leu
 145

<210> 9

<211> 1815

<212> DNA

<213> Zea mays

<220>

<223> Mlo6

<400> 9

aagcgatcga tagaataaat taaaggcgcg ggggcaacaa caagaatggg gggcggtggc 60
 ggtggcgga actcgcgga gcttgaccag acgccgacat gggcggtggc gtcggtgtgc 120
 ggcgtgatcg tgctcatctc catcctgctg gagaaggggc tccaccacgt gggcgagttc 180
 ttctcccacc gcaagaagaa ggccatggtg gaggccctgg agaaggtgaa ggcggagctc 240

```

atggtgctgg gcttcatctc gctcctcctc gtgttcggcc agaactacat catcaagggtc 300
tgcattcagca accacgccgc caacaccatg ctcccctgca agctcgaggc cgccgccgtc 360
gagggcaagg acggccacgg caaggaggcc gccgccgtgg tcgctggcaa gaagaagggtc 420
gccgtcgccg tccctggaaa gaagaagaag aaggccgccg ccgccgccga ccatcttggc 480
ggtgtggtgg actggccgcc gccctactac ggcacaaacg ccaggatgct ggcggaggcg 540
agcatggcga ccaagtgcgc cgaggggaaa gtgccgtcga tctccatcaa cgccctgcac 600
cagctgcaca tcttcatctt ctctctcgcc gtcttccacg tctctacag cgcaatcacc 660
atggcgctcg gcagggccaa gatacgtgca tggaaagagt gggagaaaga agctgcagga 720
caagactacg agttctcaca tgacccgacg cggttcaggt tcaccacga gacttcttc 780
gtgaggcagc atatgaatgt gctgaacaag ttcccagcat cattctacat cagcaacttc 840
ttccggcagt tcttcaggtc cgtgaggcag gcagactact gcgcgtgctg ccacagcttt 900
gtcaacgtcc atctggcccc tggcagcaag tttgatttcc agaagtacat caagcgggtc 960
ctggaggatg acttcaagggt gatcgtgggg atcagtcctc ctctgtgggc ttctgtctc 1020
atcttctctt tctcaacgt caatggatgg cacaccatgc tctggatctc catcatgccg 1080
gtggtgatca tctgtcgggt ggggacgaag ctgcagggca tcatctgccg catggcgatc 1140
gacatcacgg agcgccacgc cgtcatccag ggcattccga tggcgcaagt cagcgactcc 1200
tacttctggg tcgcacgccc cacttctgtg ctcttctc tccacttcac cctcttccag 1260
aatggcttcc agatcatcta ctctctctgg attctgtatg agtacggcat ggactcgtgc 1320
ttcaacgact ccgaagagtt cgtctttgca cgactctgcc ttggcgtggg tgtccagggtg 1380
ctgtgcagct acgtgacgct cccgctgtac gcgctcgtct cccaaatggg ctccaccatg 1440
aagcagtcca tcttcgacga gcagacctcc aaggcgtcga agaactggcg cgccggcgcc 1500
aagaagaagg ctcccaccgg cggctccaag cacggcgggtg gtggctcccc caccgcgggc 1560
ggcagcccca ccaaggccga cggcgacgcg tagagaggaa cacgctaact ttaatttctg 1620
tgtgcttaat tgcctaggct cgttaagtca gaacatgcat gcatgtaaca ccactgctgg 1680
ttttcatata gtgtcgacag atgggtcaacg tactttttgc gatccactt grattttttt 1740
ttacaatgaa gcacccgtcc gcgtccgtgg aactgcaag tgcattgcca aaaaaaaaaa 1800
aaaaaaaaaa aaaaaa 1815

```

<210> 10

<211> 515

<212> PRT

<213> Zea mays

<220>

<223> M106

<400> 10

Met Gly Gly Gly Gly Gly Gly Gly Asn Ser Arg Glu Leu Asp Gln Thr

1

5

10

15

Pro Thr Trp Ala Val Ala Ser Val Cys Gly Val Ile Val Leu Ile Ser

20

25

30

Ile Leu Leu Glu Lys Gly Leu His His Val Gly Glu Phe Phe Ser His

35

40

45

Arg Lys Lys Lys Ala Met Val Glu Ala Leu Glu Lys Val Lys Ala Glu

50

55

60

Leu Met Val Leu Gly Phe Ile Ser Leu Leu Leu Val Phe Gly Gln Asn
 65 70 75 80

Tyr Ile Ile Lys Val Cys Ile Ser Asn His Ala Ala Asn Thr Met Leu
 85 90 95

Pro Cys Lys Leu Glu Ala Ala Ala Val Glu Gly Lys Asp Gly His Gly
 100 105 110

Lys Glu Ala Ala Ala Val Val Ala Gly Lys Lys Lys Val Ala Val Ala
 115 120 125

Val Pro Gly Lys Lys Lys Lys Lys Ala Ala Ala Ala Ala Asp His Leu
 130 135 140

Gly Gly Val Val Asp Trp Pro Pro Pro Tyr Tyr Ala His Asn Ala Arg
 145 150 155 160

Met Leu Ala Glu Ala Ser Met Ala Thr Lys Cys Pro Glu Gly Lys Val
 165 170 175

Pro Leu Ile Ser Ile Asn Ala Leu His Gln Leu His Ile Phe Ile Phe
 180 185 190

Phe Leu Ala Val Phe His Val Ser Tyr Ser Ala Ile Thr Met Ala Leu
 195 200 205

Gly Arg Ala Lys Ile Arg Ala Trp Lys Glu Trp Glu Lys Glu Ala Ala
 210 215 220

Gly Gln Asp Tyr Glu Phe Ser His Asp Pro Thr Arg Phe Arg Phe Thr
 225 230 235 240

His Glu Thr Ser Phe Val Arg Gln His Met Asn Val Leu Asn Lys Phe
 245 250 255

Pro Ala Ser Phe Tyr Ile Ser Asn Phe Phe Arg Gln Phe Phe Arg Ser
 260 265 270

Val Arg Gln Ala Asp Tyr Cys Ala Leu Arg His Ser Phe Val Asn Val
 275 280 285

His Leu Ala Pro Gly Ser Lys Phe Asp Phe Gln Lys Tyr Ile Lys Arg
 290 295 300

Ser Leu Glu Asp Asp Phe Lys Val Ile Val Gly Ile Ser Pro Pro Leu
 305 310 315 320

Trp Ala Ser Ala Leu Ile Phe Leu Phe Leu Asn Val Asn Gly Trp His
 325 330 335

Thr Met Leu Trp Ile Ser Ile Met Pro Val Val Ile Ile Leu Ser Val
 340 345 350

Gly Thr Lys Leu Gln Gly Ile Ile Cys Arg Met Ala Ile Asp Ile Thr
 355 360 365

Glu Arg His Ala Val Ile Gln Gly Ile Pro Met Val Gln Val Ser Asp
 370 375 380

Ser Tyr Phe Trp Phe Ala Arg Pro Thr Phe Val Leu Phe Leu Ile His
 385 390 395 400

Phe Thr Leu Phe Gln Asn Gly Phe Gln Ile Ile Tyr Phe Leu Trp Ile
 405 410 415

Leu Tyr Glu Tyr Gly Met Asp Ser Cys Phe Asn Asp Ser Glu Glu Phe
 420 425 430

Val Phe Ala Arg Leu Cys Leu Gly Val Val Val Gln Val Leu Cys Ser
 435 440 445

Tyr Val Thr Leu Pro Leu Tyr Ala Leu Val Ser Gln Met Gly Ser Thr
 450 455 460

Met Lys Gln Ser Ile Phe Asp Glu Gln Thr Ser Lys Ala Leu Lys Asn
 465 470 475 480

Trp Arg Ala Gly Ala Lys Lys Lys Ala Pro Thr Gly Gly Ser Lys His
 485 490 495

Gly Gly Gly Gly Ser Pro Thr Ala Gly Gly Ser Pro Thr Lys Ala Asp
 500 505 510

Gly Asp Ala
 515

<210> 11

<211> 1660

<212> DNA

<213> Zea mays

<220>

<223> M1o7

<400> 11

```

gctagcagga gccaggtcgc gccgacagca gcagttggaa gaggatccat ggggaaggag 60
gcgacgcttg cgttcacgcc cacctgggtg gtggccatcg tatgcctcgt catcgtctct 120
atctccctcg ccgctgagcg ctactccat tacctcggca agtatctgga gtgtaagaag 180
cagaaagcgc ttttttcagc tctacagaga ctaaaagaag agctaattgct tctcggattc 240
atttcattcg tcctgagtct ctcccaaggt tttattgtta gcatttgcatt tccagaaact 300
tctactgact tcatgcttcc atgcaataga ggcaactcca gagttgcaga agaagggtgct 360
aaaatttgca aaaaaagggt tgatgttccc ttgctatcac tggaggcatt acatcagctg 420
cacattttca tatttgtact tggtttagtc catgttgtgt tctgcgctac aacaatatca 480
tttagtgagg ccaagatgag aaagtggaaag cattgggaaa cagaaattca cagagaagta 540
catgagaagt tacagcaaga aaagaatgaa ggacaagggt catcattgag tattgttgtg 600
ctacatcggg agcaccagga tgaatttgtc cacaagcggg caaagggtt ttggatgaag 660
ctcgtctgtg caagctggat aactgcattc ttgaagcaat ttcaagattc agtgagtaaa 720
tcagattatg aagcacttag atcagctttt gtcgtggaac actatcccga gaaaccagat 780
attgatttcc acaatacat gactcgcgct gttgaatatg agtttaaaaag agttgttgtg 840
atcagctggt atctgtggct ttttgaatc ttattcctgc tgctgaatat aaatggatgg 900
cacacatact tctggttggc tttcttgct ctatttctgt tacttattgt tggtgccaaa 960
ctagagcaca ttatcactcg gttggctcaa gaggcagcga tatcattatc aaataatata 1020
gaggaagtgc cgaaaataaa gccatgcaag gaccatttct ggtttcacaa gcctgagcta 1080
gtccttcatt tgatccattt catcctgttc cagaattcgt tcgagattag tttttcttc 1140
tggattctgg tatcagaagg tttcggttcg tgtatgatgg aacggaagcc ttatgtcatt 1200
tccagacttg ttatcggggg gatcatcgaa gtcactctga gctatatcac gctgccacta 1260
tacgccatcg tgaccatat gaccggacag atcaagctgc atggtttttg ctcccgctg 1320
cacgagagtg tccatggctg gatcggcttg aggaagaaac cttctcgtt ctggaagatt 1380
ccggcggtg accctaacgc tgattccggc agggaggccg acgtaaccgc cagggtggcc 1440
aaggagcgt cggggagctc gcgcagcatg ccgatggcgc cggctgacga gatcgtcacc 1500
gtcgacgacg tcgcggtggc tgcagctgcc gttggccaag gaccataaaa ttcctagcta 1560
gcaaacacca gcagggaagg caggcaggct gccgaggtag cgtgcgcagc gggttgatgg 1620
gtcggacgct gcggaaaatt cctcgtgtcg ggagcaggct 1660

```

<210> 12

<211> 499

<212> PRT

<213> Zea mays

<220>

<223> M1o7

<400> 12

```

Met Gly Lys Glu Ala Thr Leu Ala Phe Thr Pro Thr Trp Val Val Ala
  1              5              10              15

```

```

Ile Val Cys Leu Val Ile Val Ser Ile Ser Leu Ala Ala Glu Arg Ser
      20              25              30

```

```

Leu His Tyr Leu Gly Lys Tyr Leu Glu Cys Lys Lys Gln Lys Ala Leu
      35              40              45

```

Phe Ser Ala Leu Gln Arg Leu Lys Glu Glu Leu Met Leu Leu Gly Phe
 50 55 60

Ile Ser Phe Val Leu Ser Leu Ser Gln Gly Phe Ile Val Ser Ile Cys
 65 70 75 80

Ile Pro Glu Thr Ser Thr Asp Phe Met Leu Pro Cys Asn Arg Gly Asn
 85 90 95

Ser Arg Val Ala Glu Glu Gly Ala Lys Ile Cys Asn Lys Lys Gly Asp
 100 105 110

Val Pro Leu Leu Ser Leu Glu Ala Leu His Gln Leu His Ile Phe Ile
 115 120 125

Phe Val Leu Gly Leu Val His Val Val Phe Cys Ala Thr Thr Ile Ser
 130 135 140

Phe Ser Gly Ala Lys Met Arg Lys Trp Lys His Trp Glu Thr Glu Ile
 145 150 155 160

His Arg Glu Val His Glu Lys Leu Gln Gln Glu Lys Asn Glu Gly Gln
 165 170 175

Gly Ser Ser Leu Ser Ile Val Val Leu His Arg Glu His Gln Asp Glu
 180 185 190

Phe Val His Lys Arg Thr Lys Gly Phe Trp Met Lys Leu Ala Val Ala
 195 200 205

Ser Trp Ile Thr Ala Phe Leu Lys Gln Phe Gln Asp Ser Val Ser Lys
 210 215 220

Ser Asp Tyr Glu Ala Leu Arg Ser Ala Phe Val Val Glu His Tyr Pro
 225 230 235 240

Glu Lys Pro Asp Ile Asp Phe His Lys Tyr Met Thr Arg Ala Val Glu
 245 250 255

Tyr Glu Phe Lys Arg Val Val Gly Ile Ser Trp Tyr Leu Trp Leu Phe
 260 265 270

Val Ile Leu Phe Leu Leu Leu Asn Ile Asn Gly Trp His Thr Tyr Phe
 275 280 285

Trp Leu Ala Phe Leu Pro Leu Phe Leu Leu Leu Ile Val Gly Ala Lys
 290 295 300

Leu Glu His Ile Ile Thr Arg Leu Ala Gln Glu Ala Ala Ile Ser Leu
 305 310 315 320
 Ser Asn Asn Thr Glu Glu Val Pro Lys Ile Lys Pro Cys Lys Asp His
 325 330 335
 Phe Trp Phe His Lys Pro Glu Leu Val Leu His Leu Ile His Phe Ile
 340 345 350
 Leu Phe Gln Asn Ser Phe Glu Ile Ser Phe Phe Phe Trp Ile Leu Val
 355 360 365
 Ser Glu Gly Phe Gly Ser Cys Met Met Glu Arg Lys Pro Tyr Val Ile
 370 375 380
 Ser Arg Leu Val Ile Gly Val Ile Ile Glu Val Ile Cys Ser Tyr Ile
 385 390 395 400
 Thr Leu Pro Leu Tyr Ala Ile Val Thr His Met Thr Gly Gln Ile Lys
 405 410 415
 Leu His Gly Phe Gly Ser Arg Val His Glu Ser Val His Gly Trp Ile
 420 425 430
 Gly Leu Arg Lys Lys Pro Phe Ser Phe Trp Lys Ile Pro Gly Gly Asp
 435 440 445
 Pro Asn Ala Asp Ser Gly Arg Glu Ala Asp Val Thr Arg Arg Val Ala
 450 455 460
 Lys Glu Arg Ser Gly Ser Ser Arg Ser Met Pro Met Ala Pro Ala Asp
 465 470 475 480
 Glu Ile Val Thr Val Asp Asp Val Ala Val Ala Ala Ala Val Gly
 485 490 495
 Gln Gly Pro

<210> 13

<211> 1798

<212> DNA

<213> Zea mays

<220>

<223> M108

<400> 13

```

ccgggagcgg caagactggg acacgccagc cctttatttc gagctttctg cgtgcgggag 60
gggggttgac aggcgatggc ggccgagggc gaggcggcgg cgctggagtt cacaccgacg 120
tggatcgctg cggcgggtctg ctctctcatc gtgctcctct cgctcgctgc cgagcgatgc 180
ctccactacc tcggcaagac gctcaagagg aagaaccaga agccgctctt cgaggcgctg 240
ctcaagggtca aagaagagtt gatgcttctg gggttcatct ccctgctgct gacgggtgttc 300
caggggatga tccggaggac gtgcatccct gaacgctgga cattccacat gctgccatgc 360
gagaagccag atgagaaggc cggtgaggcc gccaccatgg agcattttgt agggacgctt 420
ggcaggatcg gtaggcgtct gttgcaggaa ggcactgctg gggctgagca atgccagaag 480
aagggaaaag ttccactttt gtcccttgaa gccatacatc agctgcacat tttcatattt 540
gttctggcaa tcacacatgt tattttcagc gtcacaacta tgcttttagg aggtgcacag 600
atacaccaat ggaaacagtg ggagaatgga attaaaaaag atgctcctgg aaatgggcct 660
aaggtaacca atgtacatca tcatgaattt atcaagaaac gttttaaggg tattggcaaa 720
gattctataa tattgagttg gctgcattct tttggtaagc agttttatag atcagtatct 780
aaatcagatt acaccacaat gcgtcttggg tttatcatga ctactgccc tggaaatcca 840
aaatttgatt tccatagata catggtaagg gttttagagg cggattttta gaaagtggta 900
ggcataagct ggtacttggt ggtcttcgtg gtgatatctt tgttgctgaa tgttaatggc 960
tggcacacat acttttggat tgctttccct ccccttattc ttctgttagc cattggcact 1020
aagctggagc atgtcatagc tcagctagcc catgatgtag ctgagaagca cacagcggtc 1080
gagggcgatg tgatcgtaaa accatcagat gaacacttct gggtcggcaa gcctaggggt 1140
atcctttacc tgatccactt catcctcttt cagaatgcgt ttgagattgc gtttttcttc 1200
tggaactga gcacttatgg attcgactcg tgcacatgg gacaagttcg ttttattgtg 1260
ccaaggcttg tcatcggggt gggtattcag cttctctgca gctacagcac cttgcctctg 1320
tatgcaattg taaccagat ggggagctgc tacaagaagg agatcttcaa cgagcatgtg 1380
cagcagggcg tcctgggctg ggctcagaag gtcaagatga aaaagggact gaaggagct 1440
gcatctgcta gcaaggacga atcgattacc aatgccgatt cggcaggacc ttccgttaag 1500
attgaaatgg cgaaggctgg ggaggatgtt gagatcggtt gaaacacagg ttgattggga 1560
caataggggtg cccgtgttgt aatgatgtaa cagggttaata tgccatcatc tttttttgta 1620
gatactagat agcttgctgt ggcaataccg caatagcggg gaactagaga aggtgagttt 1680
gggcccggga gcctcatctg ttatcgggtc agtaggaagc aaattcttat atacgggata 1740
tcgataagaa atgaactaag aacatgttcc tggattcatt aaaaaaaaaa aaaaaaaaaa 1798

```

<210> 14

<211> 492

<212> PRT

<213> Zea mays

<220>

<223> M108

<400> 14

```

Met Ala Ala Glu Gly Glu Ala Ala Ala Leu Glu Phe Thr Pro Thr Trp
  1                      5                      10                     15

```

```

Ile Val Ala Ala Val Cys Ser Leu Ile Val Leu Leu Ser Leu Val Ala
      20                      25                      30

```

Glu Arg Cys Leu His Tyr Leu Gly Lys Thr Leu Lys Arg Lys Asn Gln
 35 40 45
 Lys Pro Leu Phe Glu Ala Leu Leu Lys Val Lys Glu Glu Leu Met Leu
 50 55 60
 Leu Gly Phe Ile Ser Leu Leu Thr Val Phe Gln Gly Met Ile Arg
 65 70 75 80
 Arg Thr Cys Ile Pro Glu Arg Trp Thr Phe His Met Leu Pro Cys Glu
 85 90 95
 Lys Pro Asp Glu Lys Ala Gly Glu Ala Ala Thr Met Glu His Phe Val
 100 105 110
 Gly Thr Leu Gly Arg Ile Gly Arg Arg Leu Leu Gln Glu Gly Thr Ala
 115 120 125
 Gly Ala Glu Gln Cys Gln Lys Lys Gly Lys Val Pro Leu Leu Ser Leu
 130 135 140
 Glu Ala Ile His Gln Leu His Ile Phe Ile Phe Val Leu Ala Ile Thr
 145 150 155 160
 His Val Ile Phe Ser Val Thr Thr Met Leu Leu Gly Gly Ala Gln Ile
 165 170 175
 His Gln Trp Lys Gln Trp Glu Asn Gly Ile Lys Lys Asp Ala Pro Gly
 180 185 190
 Asn Gly Pro Lys Val Thr Asn Val His His His Glu Phe Ile Lys Lys
 195 200 205
 Arg Phe Lys Gly Ile Gly Lys Asp Ser Ile Ile Leu Ser Trp Leu His
 210 215 220
 Ser Phe Gly Lys Gln Phe Tyr Arg Ser Val Ser Lys Ser Asp Tyr Thr
 225 230 235 240
 Thr Met Arg Leu Gly Phe Ile Met Thr His Cys Pro Gly Asn Pro Lys
 245 250 255
 Phe Asp Phe His Arg Tyr Met Val Arg Val Leu Glu Ala Asp Phe Lys
 260 265 270
 Lys Val Val Gly Ile Ser Trp Tyr Leu Trp Val Phe Val Val Ile Phe
 275 280 285

Leu Leu Leu Asn Val Asn Gly Trp His Thr Tyr Phe Trp Ile Ala Phe
 290 295 300

Leu Pro Leu Ile Leu Leu Leu Ala Ile Gly Thr Lys Leu Glu His Val
 305 310 315 320

Ile Ala Gln Leu Ala His Asp Val Ala Glu Lys His Thr Ala Val Glu
 325 330 335

Gly Asp Val Ile Val Lys Pro Ser Asp Glu His Phe Trp Phe Gly Lys
 340 345 350

Pro Arg Val Ile Leu Tyr Leu Ile His Phe Ile Leu Phe Gln Asn Ala
 355 360 365

Phe Glu Ile Ala Phe Phe Phe Trp Ile Leu Ser Thr Tyr Gly Phe Asp
 370 375 380

Ser Cys Ile Met Gly Gln Val Arg Phe Ile Val Pro Arg Leu Val Ile
 385 390 395 400

Gly Val Val Ile Gln Leu Leu Cys Ser Tyr Ser Thr Leu Pro Leu Tyr
 405 410 415

Ala Ile Val Thr Gln Met Gly Ser Cys Tyr Lys Lys Glu Ile Phe Asn
 420 425 430

Glu His Val Gln Gln Gly Val Leu Gly Trp Ala Gln Lys Val Lys Met
 435 440 445

Lys Lys Gly Leu Lys Gly Ala Ala Ser Ala Ser Lys Asp Glu Ser Ile
 450 455 460

Thr Asn Ala Asp Ser Ala Gly Pro Ser Val Lys Ile Glu Met Ala Lys
 465 470 475 480

Ala Gly Glu Asp Val Glu Ile Val Gly Asn Thr Gly
 485 490

<210> 15

<211> 1557

<212> DNA

<213> Zea mays

<220>

<223> M109

<400> 15

cattcaccga ggagaaatct gaaaactagt taccataacc atgaaggctg ccgtgaggga 60
tatgagtcac ttgtttcgca tgaagggttg gagcagctgc accggtttat atttgtaatg 120
gctgtgactc atgtgactta tagctgcctg acgatgttac tagcaatact caagatccat 180
aaatggagaa aatgggagga tgaagcattc agggataatc atgaatcatt ttctcagatc 240
gcatatgagt cagcaactag aaggcaacca gcacttacca aatcctattc attccgctct 300
tggagccaaa ataatgtggt catgtggctt gtttgcttta ttgcgcaatt tggtcagtct 360
gttggttcgag cagactatct taccctccgc aagggtttta taatgaccca caatcttccg 420
ccaacatatg atttccacaa ttacatgata cgctcaatgg aagaggagtt tgagaagatt 480
gtcggagtgat gtggactggt gtggggcttc gttgttgctt ttatgttatt taatgtagac 540
ggatccaacc tgtacttttg gatagcaatt ctctctgtaa ctcttggtct tctgggtggg 600
gcaaagctgc agcatgtgat tgcgactcta acagcagagg gcgcaaagat gagcacctac 660
gggccaagga tacagccaag ggatgatctc ttttggttca agaaaccaga gtttctcctt 720
tggttgatac acttcgttct ctttcagaat gccttcgagt tggcttcgtt cttctggttc 780
tggtggcaat ttggttatga ttcatgcttc atcaaaaacc accttttggt atattgccgc 840
cttatactgg ggtttgctgg acagttcctg tgcagttaca gtacattacc tgtatatgcg 900
ctggtcactc agatgggggtc caagtacaag gctgcgctga tcccgaggag gatcaggag 960
accatacatg ggtggggaaa agcgacgagg aagaagcgcc ggcgccgccg cggcgacgac 1020
tcgacgggtc gcacggagac gagcacggtg tgctcgctca cggacgacga cgaagacttc 1080
gaagacgacg acgaccacca ccaccacgga ccgtcgtacg ataccccgag agccgggggc 1140
cgcccgcgtt acctgaagat cgagacgcat cgtcagtcgg gcagcgggca cgacggcccc 1200
aggcccggtg gcaccccggt cttccacccc agcggcagcg gcagcgggca cgccatgctc 1260
ctgcggcagg cctccgtctc cgcgcgctcg tcgcgctcgt acccgggagg caataacgtg 1320
acaaggtcgg cgtccatgcc cgggatcgcc gcgctgagga caacgggctc cggcacgccc 1380
acgcgcgtga gccatgagga gcctacttga tgcgcgcccc tgcatgccat ggatggcttg 1440
ctagtgcaga cgtactgagg actgagctgc tagtgtagca ccggatttga aggtatatcg 1500
atatatatat acacacgaaa agatttgagg tgacaaaaaa aaaaaaaaaa aaaaaaa 1557

<210> 16

<211> 469

<212> PRT

<213> Zea mays

<220>

<223> Mlo9

<400> 16

His Ser Pro Arg Arg Asn Leu Lys Thr Ser Tyr His Asn His Glu Gly
1 5 10 15
Cys Arg Glu Gly Tyr Glu Ser Phe Val Ser His Glu Gly Leu Glu Gln
20 25 30
Leu His Arg Phe Ile Phe Val Met Ala Val Thr His Val Thr Tyr Ser
35 40 45
Cys Leu Thr Met Leu Leu Ala Ile Leu Lys Ile His Lys Trp Arg Lys
50 55 60

Trp	Glu	Asp	Glu	Ala	Phe	Arg	Asp	Asn	His	Glu	Ser	Phe	Ser	Gln	Ile	65	70	75	80
Ala	Tyr	Glu	Ser	Ala	Thr	Arg	Arg	Gln	Pro	Ala	Leu	Thr	Lys	Ser	Tyr	85	90	95	
Ser	Phe	Arg	Ser	Trp	Ser	Gln	Asn	Asn	Val	Val	Met	Trp	Leu	Val	Cys	100	105	110	
Phe	Ile	Ala	Gln	Phe	Gly	Gln	Ser	Val	Val	Arg	Ala	Asp	Tyr	Leu	Ile	115	120	125	
Leu	Arg	Lys	Gly	Phe	Ile	Met	Thr	His	Asn	Leu	Pro	Pro	Thr	Tyr	Asp	130	135	140	
Phe	His	Asn	Tyr	Met	Ile	Arg	Ser	Met	Glu	Glu	Glu	Phe	Glu	Lys	Ile	145	150	155	160
Val	Gly	Val	Ser	Gly	Leu	Leu	Trp	Gly	Phe	Val	Val	Ala	Phe	Met	Leu	165	170	175	
Phe	Asn	Val	Asp	Gly	Ser	Asn	Leu	Tyr	Phe	Trp	Ile	Ala	Ile	Leu	Pro	180	185	190	
Val	Thr	Leu	Val	Leu	Leu	Val	Gly	Ala	Lys	Leu	Gln	His	Val	Ile	Ala	195	200	205	
Thr	Leu	Thr	Ala	Glu	Gly	Ala	Lys	Met	Ser	Thr	Tyr	Gly	Pro	Arg	Ile	210	215	220	
Gln	Pro	Arg	Asp	Asp	Leu	Phe	Trp	Phe	Lys	Lys	Pro	Glu	Phe	Leu	Leu	225	230	235	240
Trp	Leu	Ile	His	Phe	Val	Leu	Phe	Gln	Asn	Ala	Phe	Glu	Leu	Ala	Ser	245	250	255	
Phe	Phe	Trp	Phe	Trp	Trp	Gln	Phe	Gly	Tyr	Asp	Ser	Cys	Phe	Ile	Lys	260	265	270	
Asn	His	Leu	Leu	Val	Tyr	Cys	Arg	Leu	Ile	Leu	Gly	Phe	Ala	Gly	Gln	275	280	285	
Phe	Leu	Cys	Ser	Tyr	Ser	Thr	Leu	Pro	Val	Tyr	Ala	Leu	Val	Thr	Gln	290	295	300	
Met	Gly	Ser	Lys	Tyr	Lys	Ala	Ala	Leu	Ile	Pro	Arg	Arg	Ile	Arg	Glu	305	310	315	320

Thr Ile His Gly Trp Gly Lys Ala Thr Arg Lys Lys Arg Arg Arg Arg
 325 330 335

Arg Gly Asp Asp Ser Thr Val Arg Thr Glu Thr Ser Thr Val Cys Ser
 340 345 350

Leu Thr Asp Asp Asp Glu Asp Phe Glu Asp Asp Asp Asp His His His
 355 360 365

His Gly Pro Ser Tyr Asp Thr Pro Arg Ala Gly Gly Arg Pro Pro Tyr
 370 375 380

Leu Lys Ile Glu Thr His Arg Gln Ser Gly Ser Gly His Asp Gly Pro
 385 390 395 400

Arg Pro Gly Gly Thr Pro Cys Phe His Pro Ser Gly Ser Gly Ser Gly
 405 410 415

His Ala Met Leu Leu Arg Gln Ala Ser Val Ser Ala Pro Ser Ser Pro
 420 425 430

Ser Tyr Arg Gly Gly Asn Asn Val Thr Arg Ser Ala Ser Met Pro Gly
 435 440 445

Ile Ala Ala Leu Arg Thr Thr Gly Ser Gly Thr Pro Thr Arg Val Ser
 450 455 460

His Glu Glu Pro Thr
 465

<210> 17
 <211> 533
 <212> PRT
 <213> Hordeum vulgare

<400> 17
 Met Ser Asp Lys Lys Gly Val Pro Ala Arg Glu Leu Pro Glu Thr Pro
 1 5 10 15

Ser Trp Ala Val Ala Val Val Phe Ala Ala Met Val Leu Val Ser Val
 20 25 30

Leu Met Glu His Gly Leu His Lys Leu Gly His Trp Phe Gln His Arg
 35 40 45

His Lys Lys Ala Leu Trp Glu Ala Leu Glu Lys Met Lys Ala Glu Leu

50		55		60											
Met	Leu	Val	Gly	Phe	Ile	Ser	Leu	Leu	Leu	Ile	Val	Thr	Gln	Asp	Pro
65					70					75					80
Ile	Ile	Ala	Lys	Ile	Cys	Ile	Ser	Glu	Asp	Ala	Ala	Asp	Val	Met	Trp
				85					90					95	
Pro	Cys	Lys	Arg	Gly	Thr	Glu	Gly	Arg	Lys	Pro	Ser	Lys	Tyr	Val	Asp
			100					105					110		
Tyr	Cys	Pro	Glu	Gly	Lys	Val	Ala	Leu	Met	Ser	Thr	Gly	Ser	Leu	His
		115					120					125			
Gln	Leu	His	Val	Phe	Ile	Phe	Val	Leu	Ala	Val	Phe	His	Val	Thr	Tyr
	130					135					140				
Ser	Val	Ile	Thr	Ile	Ala	Leu	Ser	Arg	Leu	Lys	Met	Arg	Thr	Trp	Lys
145					150					155					160
Lys	Trp	Glu	Thr	Glu	Thr	Thr	Ser	Leu	Glu	Tyr	Gln	Phe	Ala	Asn	Asp
				165					170					175	
Pro	Ala	Arg	Phe	Arg	Phe	Thr	His	Gln	Thr	Ser	Phe	Val	Lys	Arg	His
			180					185					190		
Leu	Gly	Leu	Ser	Ser	Thr	Pro	Gly	Ile	Arg	Trp	Val	Val	Ala	Phe	Phe
	195						200					205			
Arg	Gln	Phe	Phe	Arg	Ser	Val	Thr	Lys	Val	Asp	Tyr	Leu	Thr	Leu	Arg
	210					215					220				
Ala	Gly	Phe	Ile	Asn	Ala	His	Leu	Ser	Gln	Asn	Ser	Lys	Phe	Asp	Phe
225					230					235					240
His	Lys	Tyr	Ile	Lys	Arg	Ser	Met	Glu	Asp	Asp	Phe	Lys	Val	Val	Val
				245					250					255	
Gly	Ile	Ser	Leu	Pro	Leu	Trp	Gly	Val	Ala	Ile	Leu	Thr	Leu	Phe	Leu
			260					265						270	
Asp	Ile	Asn	Gly	Val	Gly	Thr	Leu	Ile	Trp	Ile	Ser	Phe	Ile	Pro	Leu
		275					280					285			
Val	Ile	Leu	Leu	Cys	Val	Gly	Thr	Lys	Leu	Glu	Met	Ile	Ile	Met	Glu
	290					295					300				
Met	Ala	Leu	Glu	Ile	Gln	Asp	Arg	Ala	Ser	Val	Ile	Lys	Gly	Ala	Pro

[illegible]

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide primer

<400> 18

agttatccaa ggcatgccgg tggtaaaact

30

<210> 19

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
primer

<400> 19

aagtcacgga gggaggcatc acttacag

28

<210> 20

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
oligonucleotide primer

<400> 20

agagaagcca acgccawcgc ctcyatttcg tc

32

INTERNATIONAL SEARCH REPORT

Int. Application No.
PCT/US 99/15255

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07K14/415 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07K C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 04586 A (INNES JOHN CENTRE INNOV LTD ;PANSTRUGA RALPH (GB); BUESCHGES RAINE) 5 February 1998 (1998-02-05) the whole document	1-11, 13-17
X	DATABASE EMBL EMBL AC/ID AA660856, 14 November 1997 (1997-11-14) COVITZ P A ET AL.: "Expressed sequence tags from a root hair-enriched Medicago trunculata cDNA library" XP002119426 abstract	13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

19 October 1999

Date of mailing of the international search report

03/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/15255

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 99 23235 A (RAFALSKI J ANTONI ;TARAMINO GRAZIANA (US); DU PONT (US); MIAO GUO) 14 May 1999 (1999-05-14) the whole document ---	1-11, 13-17
P,X	DATABASE EMBL EMBL AC AI668283, 17 May 1999 (1999-05-17) WALBOT V: "Maize ESTs from various cDNA libraries sequenced at Stanford University" XP002119427 abstract ---	13
P,X	DATABASE EMBL EMBL AC/ID AI621523, 26 April 1999 (1999-04-26) WALBOT V: "Maize ESTs from various cDNA libraries sequenced at Stanford University" XP002119428 abstract ---	13
P,X	DATABASE EMBL EMBL AC/ID AI649550, 1 May 1999 (1999-05-01) WALBOT V: "Maize ESTs from various cDNA libraries sequenced at Stanford University" XP002119429 abstract ---	13
P,X	DATABASE EMBL EMBL AC/ID AI729043, 12 June 1999 (1999-06-12) BLEWITT M ET AL.: "ESTs from developing cotton fiber" XP002119430 abstract ---	13
P,X	DATABASE EMBL EMBL AC/ID AB011444, 4 July 1999 (1999-07-04) NEMOTO Y: "Isolation of novel early salt-responding genes from wheat (Triticum aestivum L.) by differential display" XP002119431 abstract ---	13

-/--

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 99/15255

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category ²	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>DATABASE EMBL SEQUENCE DATABASE 'Online! EMBL AC/ID AI731933, 12 June 1999 (1999-06-12) BLEWITT M ET AL.: "ESTs from developing cotton fiber" XP002119432 abstract</p> <p style="text-align: center;">-----</p>	13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/15255

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9804586 A	05-02-1998	AU 3702897 A CA 2260363 A EP 0917536 A	20-02-1998 05-02-1998 26-05-1999
WO 9923235 A	14-05-1999	AU 1450199 A	24-05-1999

